Assisted reproductive technology in Australia and New Zealand 2015

Supplementary tables

October 2017

UNSW Australia

Abbreviations and symbols ii
Introductory Notes.....

Introductory Notes

This document contains the supplementary on-line tables for the Assisted Reproductive Technology in Australia and New Zealand 2015 report. These supplementary tables should be read in conjunction with the report. A copy of the report can be found at the National Perinatal Epidemiology and Statistics Unit website:

http://npesu.unsw.edu.au/surveillance/assisted-reproductive-technology-australia-new-zealand-2015

S1:

[illegible] 3: [illegible]
[illegible] 5

[illegible]	[illegible]				
[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
Embryo transfers	52,195	621	1,911	78	552
Clinical pregnancies	16,641	191	635	25	167
Live deliveries[(b)]	13,174	153	506	20	135
Clinical pregnancies per transfer cycle (%)	31.9	30.8	33.2	32.1	30.3
Live deliveries per transfer cycle (%)	25.2	24.6	26.5	25.6	24.5

(a) Epididymal sperm is extracted by either open biopsy or percutaneous epididymal sperm aspiration (PESA).

(b) A live delivery is the delivery of one or more liveborn infants, with the birth of twins or higher order multiples counted as one live delivery.

Note: Data should be interpreted with caution due to small numbers in certain cells.

[illegible] 4: [illegible]
[illegible] 5

[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
			n		
1	8,053	13,992	23,414	1,974	47,433
2	1,416	3,470	2,563	318	7,767
•	13	121	19	4	157
[illegible]	**[illegible]**	**[illegible]**	**[illegible]**	**[illegible]**	**[illegible]**
			%		
1	84.9	79.6	90.1	86.0	85.7
2	14.9	19.7	9.9	13.9	14.0
•	0.1	0.7	0.1	0.2	0.3
[illegible]	**100**	**100**	**100**	**100**	**100**

Note: Data should be interpreted with caution due to small numbers in certain cells.

| | | |

[illegible] 7: [illegible] [illegible] **[illegible] 5**

	[illegible]		[illegible]		[illegible]	
	[illegible] (a)		[illegible] (b)		[illegible] (a)	
[illegible]	n	%	n	%	n	%
Male factor only	2,381	99.3	12	0.5	2,397	100.0
Female factor	4,620	99.1	34	0.7	4,660	100.0
Tubal disease only	611	99.8	1	0.2	612	100.0
Endometriosis only	780	99.2	5	0.6	786	100.0
Other female factor only	2,458	98.9	22	0.9	2,485	100.0
Combined female factor	771	99.2	6	0.8	777	100.0
Combined male/female factor	1,507	99.1	11	0.7	1,521	100.0
Unexplained	3,617	99.4	17	0.5	3,638	100.0
Not stated	1,863	99.1	12	0.6	1,879	100.0
[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]

(a) A live delivery is the delivery of one or more liveborn infants, with the birth of twins or higher order multiples counted as one live delivery.
(b)

Б 1 3

[illegible]

[illegible]	[illegible]			
[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
Mean (weeks)	38.2	37.2	37.0	38.1
		n		
”	144	36	1	181
28-31	117	27	1	145
32-36	1,150	331	1	1,482
< 37[(a)]	1,411	394	3	1,808
•	11,054	1,210	19	12,283
[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
		%		
”	1.2	2.2	4.5	1.3
28-31	0.9	1.7	4.5	1.0
32-36	9.2	20.6	4.5	10.5
< 37[(a)]	11.3	24.5	13.6	12.8