

**Indigenous mothers  
and their babies,  
Australia 2001–2004**

The Australian Institute of Health and Welfare (AIHW) is Australia's national health and welfare statistics and information agency. The Institute's mission is *better information and statistics for better health and wellbeing*.

The AIHW National Perinatal Statistics Unit (NPSU) is a collaborating unit of the AIHW, established in 1979. The NPSU aims at improving the health of Australian mothers and babies through the collection, analysis and reporting of information on reproductive, perinatal and maternal health. It maintains national collections on perinatal health, maternal deaths, congenital anomalies and assisted reproduction technology. The NPSU is located at

PERINATAL STATISTICS SERIES

Number 19

# **Indigenous mothers and their babies, Australia 2001–2004**

**Kate Leeds**

**Michelle Gourley**

**Paula Laws**

**Jessica Zhang**

**Fadwa Al-Yaman**

**Elizabeth A Sullivan**

**2007**

Australian Institute of Health and Welfare

Canberra

AIHW cat. no. PER 38

© Australian Institute of Health and Welfare 2007

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced without prior written permission from the Australian Institute of Health and Welfare. Requests and enquiries concerning reproduction and rights should be directed to the Head, Business Promotion and Media Unit, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601.

A complete list of the Institute's publications is available from the Business Promotion and Media Unit, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601, or via the Institute's website <[www.aihw.gov.au](http://www.aihw.gov.au)>.

ISSN 1321-8336

ISBN 978 1 74024 729 0

### **Suggested citation**

AIHW: Leeds KL, Gourley M, Laws PJ, Zhang J, Al-Yaman F & Sullivan EA 2007. Indigenous mothers and their babies, Australia 2001–2004. AIHW cat. no. PER 38. Perinatal statistics series no. 19. Canberra: AIHW.

Author affiliations: KL Leeds, PJ Laws and EA Sullivan, AIHW National Perinatal Statistics Unit; M Gourley, J Zhang and F Al-Yaman, Aboriginal and Torres Strait Islander Health and Welfare Unit, AIHW.

### **Australian Institute of Health and Welfare**

Board Chair

Hon. Peter Collins, AM, QC

Director

Penny Allbon

Any enquiries about or comments on this publication should be directed to:

AIHW National Perinatal Statistics Unit

Sydney Children's Hospital

Level 2, McNevin Dickson Building

Randwick Hospitals Campus

Randwick NSW 2031

AUSTRALIA

Phone: (02) 9382 1014

Fax: (02) 9382 1025

Email: [npsu@unsw.edu.au](mailto:npsu@unsw.edu.au)

Website: <[www.npsu.unsw.edu.au](http://www.npsu.unsw.edu.au)>

Cover image: Faye Oliver, *Summertime Bananas*, 43 x 65cm, acrylic on canvas, Irrkerlantye Arts, © artist. [info@irrkerylantyearts.com](mailto:info@irrkerylantyearts.com)

Published by the Australian Institute of Health and Welfare

Printed by

# Contents

- Acknowledgments.....vii**
- Abbreviations..... viii**
- Key findings ..... ix**
- 1 Introduction ..... 1**
  - Key maternal factors and perinatal outcomes ..... 1
  - The Perinatal National Minimum Data Set..... 2
  - Key data sources ..... 2
  - Data quality ..... 3
  - Structure of this report ..... 4
- 2 Summary data..... 5**
  - Summary ..... 5
  - Indigenous population..... 5
  - Indigenous women and births ..... 6
  - Summary measures of perinatal health..... 9
- 3 Indigenous mothers ..... 12**
  - Summary ..... 12
  - Demographics ..... 13
  - Maternal characteristics and risk factors ..... 16
  - Labour and birth characteristics ..... 27
  - Indigenous women who gave birth in hospital..... 34
- 4 Babies of Indigenous mothers ..... 37**
  - Summary ..... 37
  - Demographic profile ..... 37

The AIHW survey to assess the quality of Indigenous identification in perinatal data.....	58
Analysis of the Indigenous status of mothers.....	60
Conclusions and recommendations on trend analysis of perinatal data.....	69
Where to next.....	70
<b>6 Trends .....</b>	<b>71</b>
Summary.....	71
Background.....	71
Maternal age .....	72
Age of first-time mothers.....	74
Caesarean sections.....	76

# Acknowledgments

The Australian Institute of Health and Welfare National Perinatal Statistics Unit (NPSU) is a formally affiliated institution of the University of New South Wales (UNSW) and is linked to the School of Women's and Children's Health. We would like to acknowledge the support of NPSU by the School of Women's and Children's Health, UNSW and the Sydney Children's Hospital.

The NPSU values the time, effort and expertise contributed by all states and territories in the collection and provision of the data used in this report. We would like to acknowledge the staff members of the state and territory health authorities who provided data and reviewed the tables for this report:

- Lee Taylor and Kim Lim, Centre for Epidemiology and Research, NSW Department of Health
- Odette Taylor, Perinatal Data Collection Unit, Department of Human Services, Victoria
- Sue Cornes, Joanne Bunney and Vesna Dunne, Health Information Centre, Queensland Department of Health
- Vivien Gee and Janine Calver, Maternal and Child Health Unit, Department of Health, Western Australia
- Annabelle Chan, Kevin Priest and Joan Scott, Pregnancy Outcome Statistics Unit, Department of Health, South Australia
- Peter Mansfield and Helen Galea, Divisional Support Unit, Department of Health and Human Services, Tasmania
- Maureen Bourne and Louise Freebairn, Population Health Research Centre, ACT Health
-

# Abbreviations

ABS	Australian Bureau of Statistics
ACT	Australian Capital Territory
AHLO	Aboriginal Hospital Liaison Officer
AIHW	Australian Institute of Health and Welfare
ASGC	Australian Standard Geographical Classification
g	gram
HDSC	Health Data Standards Committee
NHDD	National Health Data Dictionary
NMDS	National Minimum Data Set
NPDC	National Perinatal Data Collection
NPSU	AIHW National Perinatal Statistics Unit
NSW	New South Wales
NT	Northern Territory
Qld	Queensland
SA	South Australia
SEIFA	Socioeconomic Indexes for Areas
Tas	Tasmania
VAED	Victorian Admitted Episodes Dataset
Vic	Victoria
WA	Western Australia
WHO	World Health Organization
n.a.	Not available
n.p.	Not published
..	Not applicable



# Key findings

This is the third report published by the AIHW on Indigenous mothers and their babies in Australia. Data from the National Perinatal Data

- The fetal death rate of babies born to Indigenous mothers declined significantly from 16 to 11 per 1,000 births, and remained stable among babies born to non-Indigenous mothers at around 6 per 1,000 births.

### **Indigenous status data quality**

Results of the assessment of the quality of Indigenous status in perinatal data in each state and territory over the period 1991–2004 showed that:

- data from New South Wales, Victoria, Queensland, Western Australia, South Australia and the Northern Territory are suitable for trends analysis from 1991 onwards
- data from the Australian Capital Territory and Tasmania are not considered stable enough to be included in trends analyses mainly because of small population size and some issues with data quality over the reporting period.

# 1 Introduction

*Indigenous mothers and their babies, Australia 2001–2004* has been prepared by the Australian Institute of Health and Welfare's (AIHW) National Perinatal Statistics Unit (NPSU) in collaboration with the AIHW's Aboriginal and Torres Strait Islander Health and Welfare Unit (ATSIHWU). It has been funded by the Office for Aboriginal and Torres Strait Islander Health (OATSIH) of the Australian Government Department of Health and Ageing.

This report follows *Indigenous mothers and their babies, Australia 1994–1996* (Day et al. 1999). It provides national information on the pregnancy and childbirth of Indigenous mothers, and the characteristics and outcomes of their babies. It also presents trends data on Indigenous mothers and their babies over the period 1991–2004. The report is based mainly on data from the National Perinatal Data Collection (NPDC). Chapter 5 of this report contains information on the ascertainment and quality of Indigenous data in the NPDC.

The purpose of *Indigenous mothers and their babies, Australia 2001–2004* is to provide Australia with epidemiological information including statistics on the Indigenous women who gave birth to liveborn or stillborn babies in 2001–2004, and on their babies. Some data on non-Indigenous women who gave birth and their babies are presented for comparison. The report also examines the quality of Indigenous data in the NPDC and changes over time in various maternal demographic characteristics and maternal and perinatal outcomes.

This report will be helpful to researchers, academics, students, policy makers and health service planners, and those providing services in reproductive health. The information presented can be used in developing policies and health services for Indigenous mothers and their babies and can assist in evaluating the impact of those policies and health service initiatives.

## Key maternal factors and perinatal outcomes

Women who identified as being Aboriginal or Torres Strait Islander represented 3.6% of women who gave birth in Australia in 2004 (Laws et al. 2006a). In general, Indigenous mothers have fewer interventions during labour and birth compared with non-Indigenous women. However, the maternal and perinatal outcomes of Indigenous mothers and their babies have consistently been shown to be poorer than those of non-Indigenous mothers. Some factors influencing these outcomes in Indigenous women who give birth include maternal age, remoteness, socioeconomic status, nutrition and smoking during pregnancy.

Indigenous mothers are, on average, 5 years younger than their non-Indigenous counterparts, and are more likely to give birth during their teenage years (Laws et al. 2006a; Powell & Dugdale 1999). In addition, Indigenous women generally have higher parity (number of previous births), are more often single, and have been shown to have poorer attendance for antenatal care (Panaretto et al. 2002).

Healthy babies are less likely to be born to Indigenous mothers who reside in remote areas, compared with women in city or regional areas (Graham et al. 2007). Approximately half of Indigenous mothers report smoking during their pregnancy and the proportion increases in more remote areas (Laws et al. 2006b). Maternal smoking during pregnancy has been shown to be associated with poor perinatal outcomes such as low birthweight, preterm birth and babies that are small for their gestational age (Chan et al. 2001).

Low birthweight, a key indicator of health status, is more common in babies born to Indigenous mothers, compared with babies born to non-Indigenous mothers (Panaretto et al. 2002; Powell & Dugdale 1999). Babies of Aboriginal mothers have been found more likely to be born preterm (Mohsin et al. 2003) and small for gestational age (DHS 2007). Reported fetal and neonatal death rates are higher in babies of Indigenous mothers compared with babies of non-Indigenous mothers (ABS & AIHW 2005).

## **The Perinatal National Minimum Data Set**

A National Minimum Data Set (NMDS) is a core set of data elements agreed to by the Statistical Information Management Committee and endorsed by the National Health Information Management Principal Committee for mandatory collection and reporting at a national level. An NMDS depends on a national agreement to collect uniform data and to supply it as part of a national collection (HDSC 2006). Definitions of all data elements included in National Minimum Data Sets are included in the AIHW's online metadata registry, METeOR.

The Perinatal NMDS is a specification for data collected on all births in Australia in hospitals, birth centres and the community. Data are collected from perinatal administrative and clinical record systems and forwarded regularly to the relevant state or territory health

## **Australian Bureau of Statistics**

The Australian Bureau of Statistics (ABS) compiles statistics and publishes reports on registrations of live births and perinatal deaths from data made available by the Registrar of Births, Deaths and Marriages in each state and territory. These data are used to compile vital statistics, and are administrative data collections that are routinely reported on year of registration rather than year of birth or year of death.

## **Data quality**

The data received from states and territories are checked for completeness, validity and

Western Australian data for Indigenous status is drawn from two sources. Western Australia has supplied Indigenous status for 2001, 2002 and 2004 mainly from their hospital morbidity system in which Indigenous status is collected using the NHDD data element. If data are missing or the mother did not give birth in hospital, Indigenous status is taken from the Notification of Case Attended form, completed by the midwife. This differs from the data source for 2003, where Indigenous status data provided to the NPSU were collected via the Western Australian perinatal form in the categories of 'Caucasian', 'Aboriginal/Torres Strait Islander' and 'Other'.

There are a small number of Aboriginal and Torres Strait Islander mothers who give birth in the Australian Capital Territory, and the proportion fluctuates from year to year, making this jurisdiction less comparable to other jurisdictions. In 2004, 54 of the 73 Aboriginal or Torres Strait Islander women who gave birth in the Australian Capital Territory were ACT residents.

## **Structure of this report**

This report is divided into six chapters:

Chapter 1 (Introduction) provides background information, describes the data sources and briefly discusses their overall limitations.

Chapter 2 (Summary data) contains summary information on the Indigenous population and Indigenous mothers and their babies. It also includes a comparison of the NPDC with birth registration data from the ABS.

Chapter 3 (Indigenous mothers) contains information on Indigenous women who gave birth in 2001–2004, including their demographic profile (e.g. maternal age), maternal characteristics (e.g. parity), and characteristics of the labour, birth and puerperium (e.g. onset of labour, method of birth).

Chapter 4 (Babies of Indigenous mothers) contains information on the characteristics and outcomes of babies born to Indigenous mothers including birth status, birthweight, gestational age, sex ratios and length of stay in hospital.

Chapter 5 (Data ascertainment and quality) contains information on the ascertainment and quality of Indigenous status data in the NPDC.

## 2 Summary data

### Summary

For 2001–2004, 35,264 women who identified as being Aboriginal or Torres Strait Islander were reported as giving birth to 35,682 babies in the National Perinatal Data Collection (NPDC) (Table 2.1). Babies born to Indigenous mothers represented 3.6% of all births in the period 2001–2004.


Live births to Indigenous mothers increased and fetal deaths decreased over the period 2001–2004. There was an average annual increase in liveborn babies born to Indigenous mothers of around 1%, and an annual average decrease in fetal deaths of around 5% (Table 2.2).

A comparison of the NPDC with the Australian Bureau of Statistics (ABS) birth registration data, which include paternal Indigenous status, showed that during the period 2001–2003, almost 5,700 more Indigenous births were identified in the ABS data collection than in the NPDC. When only live births to Indigenous mothers were considered, the NPDC contained 13% more live births (Table 2.3).

Age-standardised rate ratios for selected summary variables comparing Indigenous with non-Indigenous mothers and their babies showed that the proportion of Indigenous mothers who smoked during pregnancy was three times the proportion observed among non-Indigenous mothers. Compared with non-Indigenous mothers, Indigenous mothers had fewer instrumental vaginal deliveries in 2001–2004. The proportions of preterm birth and liveborn low birthweight babies of Indigenous mothers were twice the proportion in babies of non-Indigenous mothers (Table 2.4).

### Indigenous population

The estimated resident Aboriginal or Torres Strait Islander population in Australia was 458,500 as of 30 June 2001, or 2.4% of the total Australian population (ABS 2004). The Indigenous population has been growing over time with an average annual growth rate of





**Table 2.1: Women who gave birth and births by Indigenous status and state and territory, 2001–2004**

<b>Indigenous status</b>	<b>NSW</b>	<b>Vic</b>	<b>Qld</b>	<b>WA</b>	<b>SA</b>	<b>ACT<sup>(a)</sup></b>	<b>NT</b>	<b>Total</b>
Indigenous mothers	8,734	1,633	11,041	6,164	1,793	277	5,622	35,264
Births to Indigenous mothers	8,843	1,655	11,170	6,251	1,814	284	5,665	35,682
Live births	8,752	1,627	11,049	6,167	1,784	277	5,602	35,258
Fetal deaths	91	28	121	84	30	7	63	424
Non-Indigenous mothers	329,386	246,418	185,723	92,116	67,800	18,357	8,773	948,573
Births to non-Indigenous mothers	334,893	250,850	188,960	93,681	69,000	18,763	8,896	965,043
Live births	332,848	248,842	187,736	93,052	68,541	18,613	8,840	958,472
Fetal deaths	2,045	2,008	1,224	629	459	150	56	6,571
<b>All births<sup>(b)</sup></b>	<b>343,736</b>	<b>252,505</b>	<b>200,130</b>	<b>99,932</b>	<b>70,814</b>	<b>19,047</b>	<b>14,561</b>	<b>1,000,725</b>

(a) Care must be taken when interpreting the 2001–2004 ACT numbers as they include non-ACT-resident Indigenous mothers who gave birth in the ACT (2.6%). This is double the percentage for ACT-resident Indigenous mothers (1.3%).

(b) Does not include births to mothers whose Indigenous status was unknown.

*Notes*

1. Data for Tasmania were not available.

Live births to Indigenous mothers increased and fetal deaths decreased over the period 2001–2004 (Table 2.2). There were 8,675 live babies born to Indigenous mothers in 2001 and 8,905 in 2004, an average annual increase of around 1%. Despite a slight increase in Indigenous fetal deaths between 2002 and 2003, fetal deaths decreased by an annual average of around 5%, from 116 in 2001 to 99 in 2004.

**Table 2.2: Births to Indigenous mothers by birth status and year, 2001–2004**

Birth status	2001	2002	2003	2004	2001–2004	Average annual % change
Live births	8,675	8,827	8,851	8,905	35,258	0.9
Fetal deaths	116	102	107	99	424	–4.9
<b>All births</b>	<b>8,791</b>	<b>8,929</b>	<b>8,958</b>	<b>9,004</b>	<b>35,682</b>	<b>0.8</b>

Source: AIHW NPSU National Perinatal Data Collection 2007.

## The National Perinatal Data Collection and the Australian Bureau of Statistics birth registration data: a comparison

Table 2.3 provides a comparison between the NPDC and the ABS birth registration data. During the period 2001–2003, the NPDC reported 26,353 live births to Indigenous mothers (Table 2.3). Almost 5,700 more Indigenous births were identified in the ABS data collection than in the NPDC over this same period (32,005 births).

One explanation for this difference is that ABS birth registrations provide information on the Indigenous status of both parents whereas the NPDC includes information on the mother only. However, when comparing the NPDC data with the ABS data where the mother or both parents were identified as Indigenous, differences are still evident. There were 22,975 births to Indigenous mothers in the ABS data collection, compared with 26,353 in the NPDC, a difference of 13%. Such differences in the two data collections may also reflect the different methods and timing of the data collections. The NPDC is an epidemiological dataset collected at birth for the purpose of monitoring pregnancy. The information is usually collected by midwives, who may or may not ask a direct question regarding Indigenous status, or may obtain Indigenous status information from antenatal or hospital records. In comparison, the birth registration data is a vital statistics collection that relies on reporting by the parents or guardians, with requirements for reporting specified by individual states and territories. The differences between the two data collections are partly due to delays in the registration of, or failure to register, some live births. However, delays in registration are likely to be balanced by the late registration of births from the previous year.

Data in Table 2.3 report over the period 2001–2003 as data were not available for births that occurred in 2004 but were registered in 2005. Therefore, presenting data from 2001 to 2003 allows for the inclusion of births that occurred late in 2003 and were registered in 2004 and also includes late registrations of births in earlier years.

Of the states and territories, Queensland had the highest number of births where either or both parents were Indigenous in 2001–2003 (9,317), followed by 8,579 Indigenous births in New South Wales. The Indigenous live birth rate per 1,000 live Australian births was highest in the Northern Territory during 2001–2003 (443 per 1,000 births). This was followed by Tasmania (70 per 1,000 births) and Queensland (66 per 1,000 births). The lowest Indigenous birth rate was observed in Victoria (10 per 1,000 births) (Table 2.3).

**Table 2.3: Live births by Indigenous status and state and territory, 2001–2003**

<b>Indigenous status</b>	<b>NSW</b>	<b>Vic</b>	<b>Qld</b>	<b>WA</b>	<b>SA</b>	<b>Tas</b>	<b>ACT</b>	<b>NT</b>	<b>Total</b>
	<b>Number</b>								
<b>NPDC</b>									
Indigenous mothers	6,439	1,194	8,278	4,661	1,301	n.a.	204	4,276	26,353
<b>ABS</b>									
Both parents Indigenous	1,706	285	2,919	1,974	501	106	54	1,995	9,540

liveborn low birthweight babies of Indigenous mothers was more than double the proportion of liveborn low birthweight babies of non-Indigenous mothers. The age-standardised proportions of pret



## 3 Indigenous mothers

### Summary

During the period 2001–2004, Indigenous mothers were, on average, younger than non-Indigenous mothers, with mean ages of 24.8 years and 29.7 years respectively. Younger maternal age in Indigenous mothers was associated with residing in remote areas and socioeconomic disadvantage (Tables 3.2 and 3.3).

Almost one-third of Indigenous mothers were first-time mothers (30%), with over half of first-time Indigenous mothers aged less than 20 years (53%). Of Indigenous mothers who gave birth in 2001–2004, 16% had given birth to their fifth baby or more (Table 3.4). Of non-Indigenous mothers, 42% were having their first baby. The mean age of first-time Indigenous mothers was 20.6 years (Table 3.5) compared with 27.7 years for non-Indigenous mothers. Indigenous mothers living in regional and remote areas had more babies than those living in major cities (Table 3.6). Parity was also greater for Indigenous mothers residing in areas of greater socioeconomic disadvantage (Table 3.7).

In the five states and territories for which smoking status data were available, 51% of Indigenous mothers reported smoking during their pregnancy (Table 3.8). This was considerably higher than the crude smoking rate for non-Indigenous mothers (17%). The prevalence of smoking during pregnancy was lower in remote areas and higher in regional areas and was higher among mothers with increased socioeconomic disadvantage (Tables 3.9 and 3.10). For the three states and territories where smoking quantity data were available, of those who smoked during pregnancy, half (50%) smoked more than 10 cigarettes per day (Table 3.12). The quantity of cigarettes smoked among Indigenous mothers increased with maternal age (Table 3.13).

Indigenous mothers had fewer interventions at the onset of labour than non-Indigenous mothers, with 70% of Indigenous mothers having a spontaneous onset of labour compared with 57% of non-Indigenous mothers. Of Indigenous mothers, 19% had an induced onset of labour (Table 3.14) compared with 26% of non-Indigenous mothers. Indigenous mothers were also less likely to have instrumental vaginal deliveries (5%) and caesarean sections (22%) (Table 3.16) when compared with non-Indigenous mothers (11% and 28%, respectively). Among Indigenous mothers, spontaneous vaginal deliveries were less common in major cities and areas of greater socioeconomic advantage (Tables 3.17 and 3.18).

The majority of Indigenous women gave birth in hospital (98%) (Table 3.19). The median length of postnatal stay for these women was 3 days (Table 3.22). Of Indigenous women who gave birth in hospital, 3% were in private hospitals (Table 3.20). For non-Indigenous women, the median length of stay in hospital was 4 days and 32% gave birth in a private hospital.


# Demographics

## Maternal age

Maternal age is an important risk factor for both obstetric and perinatal outcomes, with adverse outcomes more likely to occur in younger and older mothers. In 2001–2004, 35,264 Indigenous women gave birth (Table 3.1). The mean age over this period was 24.8 years, younger than the 29.7 years for non-Indigenous mothers.

The average age of Indigenous mothers varied among the states and territories in 2001–2004. Indigenous mothers in the Australian Capital Territory were considerably older than the national average, with a mean of 26.7 years. Indigenous mothers were younger on average in the Northern Territory (23.8 years) and Western Australia (24.4 years).

There were 7,924 Indigenous mothers aged less than 20 years (23%) in 2001–2004, compared with 4% of non-Indigenous mothers (Figure 3.1). Among the states and territories, the proportion of Indigenous mothers aged less than 20 years ranged from 15% in the Australian Capital Territory to 29% in the Northern Territory.



There were 2,541 Indigenous mothers aged 35 years and over, accounting for 7% of all Indigenous mothers. This proportion was 19% in non-Indigenous mothers. The proportion of Indigenous mothers aged 35 years and over ranged from 6% in the Northern Territory and Western Australia to 9% in the Australian Capital Territory.

The overall proportion of non-Indigenous mothers in the 20–24 year age group was lower than for Indigenous mothers (14% compared with 31%). In the remaining age groups, there

was a larger proportion of non-Indigenous than Indigenous mothers (29% aged 25–29 years and 34% aged 30–34 years).

**Table 3.1: Indigenous mothers by maternal age and state and territory, 2001–2004**

Age group (years)	NSW	Vic	Qld	WA	SA	ACT	NT	Total
Mean	25.0	25.1	25.1	24.4	24.7	26.7	23.8	24.8
<b>Number</b>								
Less than 20	1,868	360	2,161	1,467	391	41	1,636	7,924
20–24	2,716	475	3,409	1,973	566	69	1,751	10,959
25–29	2,033	373	2,810	1,463	459	69	1,195	8,402
30–34	1,437	291	1,801	873	244	72	706	5,424
35 and over	672	133	860	388	133	26	329	2,541
Not stated	8	1	—	—	—	—	5	14
<b>Total</b>	<b>8,734</b>	<b>1,633</b>	<b>11,041</b>	<b>6,164</b>	<b>1,793</b>	<b>277</b>	<b>5,622</b>	<b>35,264</b>
<b>Per cent</b>								
Less than 20	21.4	22.0	19.6	23.8	21.8	14.8	29.1	22.5
20–24	31.1	29.1	30.9	32.0	31.6	24.9	31.1	31.1
25–29	23.3	22.8	25.5	23.7	25.6	24.9	21.3	23.8
30–34	16.5	17.8	16.3	14.2	13.6	26.0	12.6	15.4
35 and over	7.7	8.1	7.8	6.3	7.4	9.4	5.9	7.2
Not stated	0.1	0.1	—	—	—	—	0.1	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

*Notes*

1. Data for Tasmania were not available.
2. Data for Victoria may differ slightly from those reported in Victorian PDCU reports because of updates in the data.

Source: AIHW NPSU National Perinatal Data Collection 2007.

## Remoteness area of mother’s usual residence

Data on the geographical location of the usual residence of the mother were mapped to levels of remoteness in the Australian Standard Geographical Classification (ASGC) remoteness structure. The area of usual residence of Indigenous mothers was quite evenly spread across remoteness categories (Table 3.2). Over one in four Indigenous mothers resided in outer regional areas (28%) in 2001–2004. There were 8,321 Indigenous mothers residing in major cities (24%), and one in five (21%) lived in very remote areas. Almost one-third of Indigenous mothers lived in either remote or very remote areas. There were 5,819 Indigenous mothers (17%) residing in inner regional areas in 2001–2004.

The average age of Indigenous mothers was lower in remote (24.2 years) and very remote (24.1 years) areas when compared with major cities (25.4 years), inner regional (25.0 years) and outer regional (24.7 years) areas. Of all mothers in remote areas, 26% were aged less than 20 years compared with 20% in major cities. In contrast, the proportion of older Indigenous mothers decreased with increased remoteness—17% of Indigenous mothers in major cities were aged 30–34 years compared with 13% in very remote areas.



**Table 3.2: Indigenous mothers by maternal age and remoteness area of usual residence, 2001–2004**

Age group (years)	Major cities	Inner regional	Outer regional	Remote	Very remote	Total <sup>(a)</sup>
Mean	25.4	25.0	24.7	24.2	24.1	24.7
			<b>Number</b>			
Less than 20	1,655	1,218	2,116	1,071	1,857	7,917
20–24	2,450	1,802	3,097	1,245	2,359	10,953
25–29	2,040	1,405	2,454	870	1,627	8,396
30–34	1,440	974	1,492	580	936	5,422
35 and over	736	417	635	280	470	2,538
Not stated	—	3	3	6	2	14
<b>Total</b>	<b>8,321</b>	<b>5,819</b>	<b>9,797</b>	<b>4,052</b>	<b>7,251</b>	<b>35,240</b>
			<b>Per cent</b>			
Less than 20	19.9	20.9	21.6	26.4	25.6	22.5
20–24	29.4	31.0	31.6	30.7	32.5	31.1
25–29	24.5	24.1	25.0	21.5	22.4	23.8
30–34	17.3	16.7	15.2	14.3	12.9	15.4
35 and over	8.8	7.2	6.5	6.9	6.5	7.2
Not stated	—	0.1	—	0.1	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Excludes mothers not usually resident in Australia and those whose area of usual residence was 'Not stated'.

Source: AIHW NPSU National Perinatal Data Collection 2007.

## Socioeconomic status

Socioeconomic Indexes for Areas (SEIFA) provide a summary measure for the socioeconomic conditions within an area. The index of advantage/disadvantage was used.

More Indigenous women who gave birth were in the most disadvantaged quintile (38%) compared with other quintiles in 2001–2004, with the number of Indigenous mothers decreasing with increasing quintile of socioeconomic advantage (Table 3.3). There were only 1,160 Indigenous mothers (3%) in the least disadvantaged quintile.

Maternal age for Indigenous mothers increased with socioeconomic advantage. The average age of Indigenous mothers in the two most disadvantaged quintiles was 24.5 years, compared with 25.0 years in the third quintile, 25.5 years in the fourth quintile and 26.2 years in the fifth and least disadvantaged quintile.

**Table 3.3: Indigenous mothers by maternal age and socioeconomic status, 2001–2004**

Age group (years)	Quintile of socioeconomic disadvantage <sup>(a)</sup>					Not stated	Total
	1st quintile (most disadvantaged)	2nd quintile	3rd quintile	4th quintile	5th quintile (least disadvantaged)		
Mean	24.5	24.5	25.0	25.5	26.2	23.3	24.7
	<b>Number</b>						
Less than 20	3,138	2,278	1,371	739	213	185	7,924
20–24	4,233	3,043	2,085	1,122	284	192	10,959
25–29	3,062	2,333	1,641	963	278	125	8,402
30–34	1,909	1,450	1,062	684	253	66	5,424
35 and over	902	633	500	345	131	30	2,541
Not stated	6	4	3	—	1	—	14
<b>Total</b>	<b>13,250</b>	<b>9,741</b>	<b>6,662</b>	<b>3,853</b>	<b>1,160</b>	<b>598</b>	<b>35,264</b>
	<b>Per cent</b>						
Less than 20	23.7	23.4	20.6	19.2	18.4	30.9	22.5
20–24	31.9	31.2	31.3	29.1	24.5	32.1	31.1
25–29	23.1	24.0	24.6	25.0	24.0	20.9	23.8
30–34	14.4	14.9	15.9	17.8	21.8	11.0	15.4
35 and over	6.8	6.5	7.5	9.0	11.3	5.0	7.2
Not stated	—	—	—	—	0.1	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Socioeconomic status quintiles are determined by postcode of usual residence based on the Australian Bureau of Statistics SEIFA . The advantage/disadvantage index was used.

Source: AIHW NPSU National Perinatal Data Collection 2007.

## Maternal characteristics and risk factors

### Parity

Parity is the number of previous pregnancies that resulted in live births or stillbirths.

Primiparous women are those who have had no previous pregnancy resulting in a live birth or stillbirth and multiparous women are those who have had at least one previous pregnancy resulting in a live birth or stillbirth.

During the 2001–2004 period, there were 10,608 Indigenous mothers who gave birth to their first baby (30%), 8,455 (24%) who gave birth to their second baby, and 6,227 (18%) who gave birth for the third time (Table 3.4). Only 12% of Indigenous women gave birth to their fourth baby, but 16% gave birth for the fifth (or more) time. The figures for non-Indigenous women who gave birth were 42% having their first baby, 34% their second, 15% their third, 5% their fourth and only 3% gave birth for the fifth time or more.

Indigenous mothers in Western Australia were more likely than mothers in others states and territories to have a parity of three or more. In Western Australia, 13% of Indigenous mothers had given birth three times previously and 19% four or more times, compared with 12% and 16% respectively for Australia.

**Table 3.4: Indigenous mothers by parity and state and territory, 2001–2004**

Parity	NSW	Vic	Qld	WA	SA	ACT	NT	Total
<b>Number</b>								
None	2,725	585	3,155	1,720	568	106	1,749	10,608
One	2,142	436	2,558	1,379	437	76	1,427	8,455
Two	1,524	263	1,992	1,090	295	34	1,029	6,227
Three	974	164	1,362	794	217	26	646	4,183
Four or more	1,339	185	1,974	1,181	276	35	765	5,755
Not stated	30	—	—	—	—	—	6	36
<b>Total</b>	<b>8,734</b>	<b>1,633</b>	<b>11,041</b>	<b>6,164</b>	<b>1,793</b>	<b>277</b>	<b>5,622</b>	<b>35,264</b>
<b>Per cent</b>								
None	31.2	35.8	28.6	27.9	31.7	38.3	31.1	30.1
One	24.5	26.7	23.2	22.4	24.4	27.4	25.4	24.0
Two	17.4	16.1	18.0	17.7	16.5	12.3	18.3	17.7
Three	11.2	10.0	12.3	12.9	12.1	9.4	11.5	11.9
Four or more	15.3	11.3	17.9	19.2	15.4	12.6	13.6	16.3
Not stated	0.3	—	—	—	—	—	0.1	0.1
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

*Notes*

1. Data for Tasmania were not available.
2. Data for Victoria may differ slightly from those reported in Victorian PDCU reports because of updates in the data.

Source: AIHW NPSU National Perinatal Data Collection 2007.


The average age of first-time Indigenous mothers was 20.6 years in 2001–2004 (Table 3.5). This was considerably younger than the average age of non-Indigenous first-time mothers (27.7 years). The average age of Indigenous mothers rose with higher parity. For each additional parity the average age increased approximately 2 to 3 years.

**Table 3.5: Indigenous mothers by maternal age and parity, 2001–2004**

Age group (years)	Parity						Total
	None	One	Two	Three	Four or more	Not stated	
Mean	20.6	23.5	25.8	27.7	30.7	30.6	24.7
	<b>Number</b>						
Less than 20	5,662	1,890	330	33	4	5	7,924
20–24	3,099	3,739	2,552	1,108	458	3	10,959
25–29	1,067	1,659	2,005	1,698	1,969	4	8,402
30–34	568	853	957	954	2,078	14	5,424
35 and over	207	312	380	387	1,245	10	2,541
Not stated	5	2	3	3	1	—	14
<b>Total</b>	<b>10,608</b>	<b>8,455</b>	<b>6,227</b>	<b>4,183</b>	<b>5,755</b>	<b>36</b>	<b>35,264</b>
	<b>Per cent</b>						
Less than 20	53.4	22.4	5.3	0.8	0.1	13.9	22.5
20–24	29.2	44.2	41.0	26.5	8.0	8.3	31.1
25–29	10.1	19.6	32.2	40.6	34.2	11.1	23.8
30–34	5.4	10.1	15.4	22.8	36.1	38.9	15.4
35 and over	2.0	3.7	6.1	9.3	21.6	27.8	7.2
Not stated	—	—	—	0.1	—	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Source: AIHW NPSU National Perinatal Data Collection 2007.

Over half of first-time Indigenous mothers were aged less than 20 years (53%) (Table 3.5; Figure 3.2). Nine percent of multiparous Indigenous mothers were also in this age group. The proportion of primiparous Indigenous mothers decreased with advancing age, to 5% for



Source: AIHW NPSU National Perinatal Data Collection 2007.

**Figure 3.2: Primiparous and multiparous Indigenous mothers by maternal age, 2001-2004 (per cent)**



**Table 3.7: Indigenous mothers by parity and socioeconomic status, 2001–2004**

---

Parity	Quintile of socioeconomic disadvantage <sup>(a)</sup>	
	1st quintile (most disadvantaged)	2nd

When only stated responses are considered for all five states and territories, the percentage of Indigenous mothers recorded as smoking during pregnancy was 54%. For non-Indigenous mothers, the comparable percentage remained at 17%.

**Table 3.8: Indigenous mothers by tobacco smoking status during pregnancy and state and territory, 2001–2004**

Smoking status	NSW	WA	SA <sup>(a)</sup>	ACT	NT <sup>(b)</sup>	Total
<b>Number</b>						
Smoked	5,028	3,037	1,056	126	2,249	11,496
Did not smoke	3,697	3,127	649	150	2,370	9,993
Not stated	9	—	88	1	1,003	1,101
<b>Total</b>	<b>8,734</b>	<b>6,164</b>	<b>1,793</b>	<b>277</b>	<b>5,622</b>	<b>22,590</b>
<b>Per cent</b>						
Smoked <sup>(c)</sup>	57.6	49.3	58.9	45.5	40.0	50.9
Did not smoke	42.3	50.7	36.2	54.2	42.2	44.2
Not stated	0.1	—	4.9	0.4	17.8	4.9
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) For South Australia, 'Smoked' includes women who quit before the first antenatal visit.

(b) For the Northern Territory, smoking status was recorded at the first antenatal visit.

(c) If the Not stated responses are proportioned out, the percentage of Indigenous mothers who smoked during pregnancy was 54% (NSW 57.6%, WA 49.3%, SA 58.9%, ACT 45.5%, NT 40.0%, Total 50.9%).



**Table 3.9: Indigenous mothers by tobacco smoking status during pregnancy and remoteness area of usual residence, 2001–2004**

Smoking status	Major cities	Inner regional	Outer regional	Remote	Very remote	Total <sup>(a)</sup>
						<b>Number</b>
Smoked	3,011	1,965	2,639	1,459	2,411	11,485
Did not smoke	2,832	1,542	1,935	1,326	2,350	9,985
Not stated	23	7	196	220	651	1,097
<b>Total</b>	<b>6,867</b>	<b>3,521</b>	<b>4,770</b>	<b>3,005</b>	<b>5,412</b>	<b>20,575</b>

The proportion of Indigenous mothers who smoked during pregnancy in 2001–2004 generally increased with increased socioeconomic disadvantage (Table 3.10). The highest proportion of Indigenous mothers who smoked was in the most disadvantaged quintile (55%) and the lowest proportion was in the least disadvantaged quintile (43%). Interestingly, smoking rates for Indigenous mothers were higher in the third most disadvantaged quintile (54%), compared with rates in the second most disadvantaged quintile (47%). Forty-six percent of Indigenous mothers in the fourth quintile smoked during pregnancy.

**Table 3.10: Indigenous mothers by tobacco smoking status during pregnancy and socioeconomic status, 2001–2004**

Smoking status	Quintile of socioeconomic disadvantage <sup>(a)</sup>						Total
	1st quintile (most disadvantaged)	2nd quintile	3rd quintile	4th quintile	5th quintile (least disadvantaged)	Not stated	
<b>Number</b>							
Smoked	4,825	2,825	1,927	1,368	380	171	11,496
Did not smoke	3,464	2,852	1,557	1,451	480	189	9,993
Not stated	488	292	100	127	20	74	1,101
<b>Total<sup>(b)</sup></b>	<b>8,777</b>	<b>5,969</b>	<b>3,584</b>	<b>2,946</b>	<b>880</b>	<b>434</b>	<b>22,590</b>
<b>Per cent</b>							
Smoked	55.0	47.3	53.8	46.4	43.2	39.4	50.9
Did not smoke	39.5	47.8	43.4	49.3	54.5	43.5	44.2
Not stated	5.6	4.9	2.8	4.3	2.3	17.1	4.9
<b>Total<sup>(b)</sup></b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Socioeconomic status quintiles are determined by postcode of usual residence based on the Australian Bureau of statistics SEIFA. The advantage/disadvantage index was used.

(b) Data on tobacco smoking during pregnancy were not available for Victoria, Queensland or Tasmania.

Source: AIHW NPSU National Perinatal Data Collection 2007.

Smoking during pregnancy varied little across age groups for Indigenous mothers. The percentage of smokers reported ranged from 49% in those aged less than 20 years to 52% in those aged 20–24 years (Table 3.11).

**Table 3.11 Indigenous mothers by tobacco smoking status during pregnancy and maternal age, 2001–2004**

Smoking status	Age group (years)						Total
	Less than 20	20–24	25–29	30–34	35 and over	Not stated	
	<b>Number</b>						
Smoked	2,671	3,669	2,684	1,683	781	8	11,496
Did not smoke	2,399	3,044	2,316	1,526	704	4	9,993
Not stated	333	362	219	123	63	1	1,101
<b>Total<sup>(a)</sup></b>	<b>5,403</b>	<b>7,075</b>	<b>5,219</b>	<b>3,332</b>	<b>1,548</b>	<b>13</b>	<b>22,590</b>
	<b>Per cent</b>						
Smoked	49.4	51.9	51.4	50.5	50.5	61.5	50.9
Did not smoke	44.4	43.0	44.4	45.8	45.5	30.8	44.2
Not stated	6.2	5.1	4.2	3.7	4.1	7.7	4.9
<b>Total<sup>(a)</sup></b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Data on tobacco smoking during pregnancy were not available for Victoria, Queensland or Tasmania.

Source: AIHW NPSU National Perinatal Data Collection 2007.

Data on the average number of cigarettes smoked per day in the second half of pregnancy

**Table 3.13: Indigenous mothers who smoked during the second half of pregnancy by average number of cigarettes per day and maternal age, 2001–2004**

Average no. cigarettes per day	Age group (years)					Not stated	Total
	Less than 20	20–24	25–29	30–34	35 and over		
	<b>Number</b>						
None	46	45	30	20	6	—	147
10 or less	707	913	551	336	140	—	2,647
More than 10	576	910	797	537	308	4	3,132
Not stated	51	94	58	51	29	1	284
<b>Total<sup>(a)</sup></b>	<b>1,380</b>	<b>1,962</b>	<b>1,436</b>	<b>944</b>	<b>483</b>	<b>5</b>	<b>6,210</b>
	<b>Per cent</b>						
None	3.3	2.3	2.1	2.1	1.2	—	2.4
10 or less	51.2	46.5	38.4	35.6	29.0	—	42.6
More than 10	41.7	46.4	55.5	56.9	63.8	80.0	50.4
Not stated	3.7	4.8	4.0	5.4	6.0	20.0	4.6
<b>Total<sup>(a)</sup></b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Data on the average daily number of cigarettes smoked during the second half of pregnancy were available only in New South Wales, South Australia and the Australian Capital Territory.

Source: AIHW NPSU National Perinatal Data Collection 2007.

## Labour and birth characteristics

### Onset of labour

Onset of labour is defined as spontaneous, induced, or no labour. When examining the rates of spontaneous, induced and no labour among Indigenous mothers it becomes evident that, when compared with non-Indigenous mothers, Indigenous mothers have fewer interventions at the onset of labour (induction) and during labour (augmentation).

On 2001-2004, 70% of Indigenous women (91%) who smoked during the second half of pregnancy had a spontaneous onset of labour, compared with 79% of non-Indigenous women (81%).



## Presentation at birth

Data are included in this section by mother; for multiple births, the presentation at birth of the firstborn baby is used.

In 2001–2004, the predominant presentation at birth was vertex, occurring in 95% of Indigenous women who gave birth (Table 3.15). The proportion of vertex presentations was the same for non-Indigenous mothers during the same period (95%). Breech presentations occurred in 4% of Indigenous women who gave birth. The proportion varied across the states and territories from 3% in Victoria to 5% in the Northern Territory and South Australia. Breech presentations occurred in 5% of non-Indigenous mothers. Face and brow presentations occurred in 0.3% and other presentations in 0.8% of Indigenous women who gave birth.

**Table 3.15: Indigenous mothers by presentation of birth and state and territory, 2001–2004**

Presentation	NSW	Vic	Qld	WA	SA	ACT	NT	Total
<b>Number</b>								
Vertex	8,293	1,558	10,490	5,856	1,679	265	5,285	33,426
Breech	344	47	437	255	82	10	259	1,434
Face and brow	22	6	17	23	6	—	15	89
Other	71	18	94	30	22	2	44	281
Not stated	4	4	3	—	4	—	19	34
<b>Total</b>	<b>8,734</b>	<b>1,633</b>	<b>11,041</b>	<b>6,164</b>	<b>1,793</b>	<b>277</b>	<b>5,622</b>	<b>35,264</b>
<b>Per cent</b>								
Vertex	95.0	95.4	95.0	95.0	93.6	95.7	94.0	94.8
Breech	3.9	2.9	4.0	4.1	4.6	3.6	4.6	4.1
Face and brow	0.3	0.4	0.2	0.4	0.3	—	0.3	0.3
Other	0.8	1.1	0.9	0.5	1.2	0.7	0.8	0.8
Not stated	—	0.2	—	—	0.2	—	0.3	0.1
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

### Notes


1. Data for Tasmania were not available.
2. For multiple births, the presentation of the firstborn baby was used.

Source: AIHW NPSU National Perinatal Data Collection 2007.

## Method of birth

Data are presented in this section by mother; for multiple births, the method of birth of the firstborn baby is presented.

Of all Indigenous women who gave birth in 2001–2004, 72% had a spontaneous vaginal birth (Table 3.16). This was higher than the proportion of non-Indigenous mothers who had a spontaneous vaginal birth (61%). Compared with non-Indigenous mothers, Indigenous mothers were less likely to have forceps or vacuum extraction deliveries (11% compared with 5%) and more likely to have a vaginal breech birth (0.8% compared with 0.4%) (Figure 3.4). The caesarean section rate for Indigenous mothers (22%) was less than the rate for non-Indigenous mothers (28%).





**Table 3.16: Indigenous mothers by method of birth and state and territory, 2001–2004**

Method of birth	NSW	Vic	Qld	WA	SA	ACT	NT	Total
<b>Number</b>								
Spontaneous vaginal	6,392	1,150	8,059	4,488	1,143	182	3,932	25,346
Forceps	164	59	119	95	44	13	87	581
Vacuum extraction	275	61	275	242	60	19	183	1,115
Vaginal breech	55	16	90	67	22	3	45	298
Caesarean section	1,848	346	2,491	1,272	524	60	1,367	7,908
Other	—	—	7	—	—	—	—	7
Not stated	—	1	—	—	—	—	8	9
<b>Total</b>	<b>8,734</b>	<b>1,633</b>	<b>11,041</b>	<b>6,164</b>	<b>1,793</b>	<b>277</b>	<b>5,622</b>	<b>35,264</b>
<b>Per cent</b>								
Spontaneous vaginal	73.2	70.4	73.0	72.8	63.7	65.7	69.9	71.9
Forceps	1.9	3.6	1.1	1.5	2.5	4.7	1.5	1.6
Vacuum extraction	3.1	3.7	2.5	3.9	3.3	6.9	3.3	3.2
Vaginal breech	0.6	1.0	0.8	1.1	1.2	1.1	0.8	0.8
Caesarean section	21.2	21.2	22.6	20.6	29.2	21.7	24.3	22.4
Other	—	—	0.1	—	—	—	—	—
Not stated	—	0.1	—	—	—	—	0.1	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

*Notes*

1. Data for Tasmania were not available.
2. For multiple births, the method of birth of the firstborn baby was used.

Source: AIHW NPSU National Perinatal Data Collection 2007.

The method of birth for Indigenous mothers varied across remoteness areas within Australia. Spontaneous vaginal births were less common in major cities and remote areas (both 71%) when compared with regional and very remote areas; assisted vaginal births were more common in major cities (Table 3.17). Vacuum extraction was the method of birth for approximately 4% of Indigenous mothers who lived in major cities, compared to around 3% in remote and very remote areas.

The proportion of caesarean sections among Indigenous mothers did not vary greatly by remoteness area (between 22% and 23%).

**Table 3.17: Indigenous mothers by method of birth and remoteness area of usual residence, 2001–2004**

Method of birth	Major cities	Inner regional	Outer regional	Remote	Very remote	Total <sup>(a)</sup>
			<b>Number</b>			
Spontaneous vaginal	5,865	4,197	7,152	2,890	5,223	25,327
Forceps	158	124	146	76	77	581
Vacuum extraction	346	173	287	103	205	1,114



**Table 3.19: Indigenous mothers by actual place of birth and state and territory, 2001–2004**

Place of birth	NSW	Vic	Qld	WA	SA	ACT	NT	Total
<b>Number</b>								
Hospital	8,555	1,595	10,902	6,064	1,719	268	5,358	34,461
Birth centre	80	n.p.	21	8	54	<5	—	178
Home	4	<4	8	9	<4	—	11	35
Other	95	26	110	83	n.p.	n.p.	<sup>(a)</sup> 253	590
Not stated	—	—	—	—	—	—	—	—
<b>Total</b>	<b>8,734</b>	<b>1,633</b>	<b>11,041</b>	<b>6,164</b>	<b>1,793</b>	<b>277</b>	<b>5,622</b>	<b>35,264</b>
<b>Per cent</b>								
Hospital	98.0	97.7	98.7	98.4	95.9	96.8	95.3	97.7
Birth centre	0.9	n.p.	0.2	0.1	3.0	n.p.	—	0.5
Home	—	n.p.	0.1	0.1	n.p.	—	0.2	0.1
Other	1.1	1.6	1.0	1.3	n.p.	n.p.	<sup>(a)</sup> 4.5	1.7
Not stated	—	—	—	—	—	—	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) The majority of these births occurred in remote community health centres.

*Notes*

1. Data for Tasmania were not available.
  2. For multiple births, the place of birth of the firstborn baby was used.
- n.p. Data not published to maintain confidentiality of small numbers.

Source: AIHW NPSU National Perinatal Data Collection 2007.

## Indigenous women who gave birth in hospital

### Hospital sector

‘Hospital sector’ indicates whether a patient was admitted to a public or private hospital. Of Indigenous mothers who gave birth in hospital in 2001–2004, the proportion in private hospitals was approximately 3% (Table 3.20). This was substantially lower than the proportion of non-Indigenous mothers who gave birth in private hospitals (32%).

There was variation in the proportion of Indigenous women who gave birth in private hospitals among the states and territories; ranging from 1.1% in the Northern Territory to 11% in the Australian Capital Territory. Indigenous women who gave birth in private hospitals were older, with a mean age of 29.0 years compared with 24.6 years for Indigenous women who gave birth in public hospitals.

**Table 3.20: Indigenous mothers who gave birth in hospital by hospital sector, 2001–2004**

---

Hospital sector	NSW	Vic	Qld	WA	SA	ACT	NT	Total
-----------------	-----	-----	-----	----	----	-----	----	-------

---

## Postnatal length of stay

The median length of stay in hospital for Indigenous women who gave birth in 2001–2004 was 3 days overall and for all age groups (Table 3.22). The median length of stay for non-Indigenous mothers who gave birth over this same period was 4 days.

Almost one-quarter of Indigenous mothers who gave birth in hospital had a length of stay of 2 days (23%) and one-fifth stayed for 3 days (21%). Older mothers were generally more likely to stay in hospital for longer periods after giving birth.

**Table 3.22: Indigenous mothers who gave birth in hospital<sup>(a)</sup> by postnatal length of stay and maternal age, 2001–2004**

Length of stay	Age group (years)					Not stated	Total
	Less than 20	20–24	25–29	30–34	35 and over		
Median length of stay	3.0	3.0	3.0	3.0	3.0	4.5	3.0
	<b>Number</b>						
Less than 1 day	156	364	270	132	66	1	989
1 day	871	1,544	1,281	702	283	—	4,681
2 days	1,310	2,048	1,461	913	389	—	6,121
3 days	1,278	1,715	1,252	822	365	—	5,432
4 days	847	1,076	808	632	304	—	3,667
5 days	560	611	549	443	219	—	2,382
6 days	268	288	298	204	126	—	1,184
7–13 days	412	411	326	245	160	1	1,555
14–20 days	26	27	16	20	23	—	112
21–27 days	2	2	5	3	1	—	13
28 or more days	1	1	4	3	2	—	11
Not stated	68	115	84	61	31	5	364
<b>Total</b>	<b>5,799</b>	<b>8,202</b>	<b>6,354</b>	<b>4,180</b>	<b>1,969</b>	<b>7</b>	<b>26,511</b>
	<b>Per cent</b>						
Less than 1 day	2.7	4.4	4.2	3.2	3.4	14.3	3.7
1 day	15.0	18.8	20.2	16.8	14.4	—	17.7
2 days	22.6	25.0	23.0	21.8	19.8	—	23.1
3 days	22.0	20.9	19.7	19.7	18.5	—	20.5
4 days	14.6	13.1	12.7	15.1	15.4	—	13.8
5 days	9.7	7.4	8.6	10.6	11.1	—	9.0
6 days	4.6	3.5	4.7	4.9	6.4	—	4.5
7–13 days	7.1	5.0	5.1	5.9	8.1	14.3	5.9
14–20 days	0.4	0.3	0.3	0.5	1.2	—	0.4
21–27 days	—	—	0.1	0.1	0.1	—	—
28 or more days	—	—	0.1	0.1	0.1	—	—
Not stated	1.2	1.4	1.3	1.5	1.6	71.4	1.4
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

## 4 Babies of Indigenous mothers

### Summary

The perinatal outcomes of babies born to Indigenous mothers were poor when compared with babies born to non-Indigenous mothers during 2001–2004. Fourteen per cent of births to Indigenous mothers were preterm (Table 4.2) compared with 8% of babies born to non-Indigenous mothers. Older Indigenous mothers were more likely to give birth to a preterm baby (Table 4.4).

Thirteen per cent of liveborn babies of Indigenous mothers were low birthweight over the 2001–2004 period (Table 4.5). This was more than double the proportion observed among liveborn babies of non-Indigenous mothers (6%). The proportion of low birthweight babies among babies born to Indigenous mothers was highest in older mothers (15% in those aged 35 years and over) (Table 4.8). The mean birthweight of babies born to Indigenous mothers decreased with increased remoteness and socioeconomic disadvantage (Tables 4.6 and 4.7).

Of liveborn babies of Indigenous mothers, approximately 3% had a low Apgar score at 5 minutes (Table 4.10). This was higher than the proportion observed among liveborn babies of non-Indigenous mothers (1.3%). The proportion of babies born to Indigenous mothers who had a low Apgar score at 5 minutes increased with increased remoteness (Table 4.11).

The median length of stay in hospital for babies of Indigenous mothers was 3 days (Table 4.12), shorter than the median length of stay observed among babies born to non-Indigenous mothers (4 days). The median length of stay among babies born to Indigenous mothers was higher in very remote areas and in the most advantaged socioeconomic quintile (both 4 days) (Tables 4.13 and 4.14).

There were 424 fetal deaths and 201 neonatal deaths reported among babies born to Indigenous mothers for 2001–2004. This equated to a fetal death rate of 12 per 1,000 births and a neonatal death rate of 6 per 1,000 live births (Table 4.15). This was considerably higher than the fetal and neonatal death rates observed among babies born to non-Indigenous mothers (7 and 3 per 1,000 births respectively).

### Demographic profile

#### Birth status

Babies are recorded as liveborn or stillborn (fetal deaths) on perinatal notification forms. A live birth is defined by the World Health Organisation (WHO) as the complete expulsion or extraction from the mother of a baby which, after such separation, breathes or shows any other evidence of life. A fetal death is defined as a death occurring before the complete expulsion or extraction from the mother of a product of conception of 20 or more completed weeks gestation or 400 grams or more birthweight (NHDC 2003). The same criteria are applied to live births for inclusion in the NPDC; that is, live births must also be at least 20 weeks gestation or at least 400 grams birthweight.

There were 35,258 live births and 424 fetal deaths to Indigenous mothers in Australia in 2001–2004, giving a total of 35,682 births to Indigenous mothers reported to the NPDC (Table 2.1).

## Sex

Male live births to Indigenous mothers exceeded female live births in all states and territories, except Victoria and the Australian Capital Territory, and accounted for 51% of live births to Indigenous mothers in 2001–2004 (Table 4.1). The sex ratio (defined as the number of male liveborn babies per 100 female liveborn babies) for babies born to Indigenous mothers was 105 nationally and ranged from 95 in the Australian Capital Territory to 109 in the Northern Territory. For live births to non-Indigenous mothers, the sex ratio was 106 males per 100 females.

**Table 4.1: Live births to Indigenous mothers by sex and state and territory, 2001–2004**

Sex of baby	NSW	Vic	Qld	WA	SA	ACT	NT	Total
Sex ratio (M:F)	104.4	96.3	107.9	100.7	101.7	95.1	108.8	104.9
	<b>Number</b>							
Males	4,469	798	5,735	3,094	899	135	2,918	18,048
Females	4,281	829	5,314	3,073	884	142	2,683	17,206
Indeterminate/ not stated	2	—	—	—	1	—	1	4
<b>Total</b>	<b>8,752</b>	<b>1,627</b>	<b>11,049</b>	<b>6,167</b>	<b>1,784</b>	<b>277</b>	<b>5,602</b>	<b>35,258</b>
	<b>Per cent</b>							
Males	51.1	49.0	51.9	50.2	50.4	48.7	52.1	51.2
Females	48.9	51.0	48.1	49.8	49.6	51.3	47.9	48.8
Indeterminate/ not stated	—	—	—	—	0.1	—	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Note: Data for Tasmania were not available.

Source: AIHW NPSU National Perinatal Data Collection 2007.

## Outcomes


### Gestational age

Preterm birth (before 37 weeks gestation) is associated with neonatal problems that cause significant morbidity and mortality in newborn babies. Preterm births were classified according to the criteria of the WHO into groups of 20–27 weeks, 28–31 weeks and 32–36 weeks.

In 2001–2004 there were 4,962 preterm births to Indigenous mothers, accounting for 14% of all births to Indigenous mothers (Table 4.2). This was almost double the proportion of preterm births to non-Indigenous mothers (8%) (Figure 4.1). There was variation in the proportion of preterm births to Indigenous mothers among the states and territories, ranging



from 12% of all births in New South Wales to 22% in the Australian Capital Territory. A large proportion of Indigenous women who gave birth in the Australian Capital Territory were residents of New South Wales, therefore, preterm births appear high when based on births to Indigenous women who gave birth in the Australian Capital Territory. Note that there are a small number of Aboriginal and Torres Strait Islander mothers who give birth in the Australian Capital Territory, and the proportion fluctuates from year to year, making this jurisdiction less comparable to other jurisdictions.



Note: For multiple births the gestational age of the firstborn baby was used for all babies.

Source: AIHW NPSU National Perinatal Data Collection 2007.

**Figure 4.1: Gestational age: Indigenous and Non-Indigenous babies**

**Table 4.3: Births to Indigenous mothers by gestational age and plurality, 2001–2004**

Gestational age (weeks)	Plurality			Total
	One	Two	Three or more	
	<b>Number</b>			
20–27 <sup>(a)</sup>	555	64	6	625
28–31	511	76	9	596
32–36	3,381	354	6	3,741
37–41	29,826	316	—	30,142
42 and over	560	—	—	560
Not stated	18	—	—	18
<b>Total</b>	<b>34,851</b>	<b>810</b>	<b>21</b>	<b>35,682</b>
20–36 <sup>(a)</sup>	4,447	494	21	4,962
	<b>Per cent</b>			
20–27 <sup>(a)</sup>	1.6	7.9	28.6	1.8
28–31	1.5	9.4	42.9	1.7
32–36	9.7	43.7	28.6	10.5
37–41	85.6	39.0	—	84.5
42 and over	1.6	—	—	1.6
Not stated	0.1	—	—	0.1
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
20–36 <sup>(a)</sup>	12.8	61.0	100.0	13.9

(a) Includes two babies of less than 20 weeks gestation.

Note: For multiple births the gestational age of the firstborn baby was used for all babies.

Source: AIHW NPSU National Perinatal Data Collection 2007.



## Birthweight

A baby's birthweight is a key indicator of health status. Babies are defined as low birthweight if their weight at birth is less than 2,500 grams. Within this category, those weighing less than 1,500 grams are defined as very low birthweight (WHO 1992).

In 2001–2004, 86% of liveborn babies to Indigenous mothers had a birthweight in the range of 2,500–4,499 grams (Table 4.5). The mean birthweight was 3,162 grams. Thirteen per cent of liveborn babies to Indigenous mothers fell within the low birthweight range. This proportion was considerably higher than the proportion of liveborn babies to non-Indigenous mothers who had a birthweight of less than 2,500 grams (6%). The mean birthweight of liveborn babies of non-Indigenous mothers was 3,381 grams.

The proportion of low birthweight liveborn babies born to Indigenous mothers varied among the states and territories; from 12% in both New South Wales and Queensland to 19% in the Australian Capital Territory. A large proportion of Indigenous women who gave birth in the Australian Capital Territory were residents of New South Wales, therefore, the proportion of low birthweight babies appears high when based on births to Indigenous women who gave birth in the Australian Capital Territory.

**Table 4.5: Live births to Indigenous mothers by birthweight and state and territory, 2001–2004**

Birthweight (g)	NSW	Vic	Qld	WA	SA	ACT <sup>(a)</sup>	NT	Total
Mean	3,193	3,158	3,200	3,124	3,064	3,085	3,117	3,162
	<b>Number</b>							
Less than 1,500	165	38	255	160	78	18	127	841
1,500 – 2,499	903	184	1,020	732	236	35	627	3,737
2,500 – 4,499	7,560	1,375	9,586	5,199	1,445	223	4,781	30,169
4,500 and over	120	29	185	75	n.p.	<5	66	501
Not stated	4	1	3	1	n.p.	n.p.	1	10
<b>Total</b>	<b>8,752</b>	<b>1,627</b>	<b>11,049</b>	<b>6,167</b>	<b>1,784</b>	<b>277</b>	<b>5,602</b>	<b>35,258</b>
<i>Less than 2,500</i>	<i>1,068</i>	<i>222</i>	<i>1,275</i>	<i>892</i>	<i>314</i>	<i>53</i>	<i>754</i>	<i>4,578</i>
	<b>Per cent</b>							
Less than 1,500	1.9	2.3	2.3	2.6	4.4	6.5	2.3	2.4
1,500 – 2,499	10.3	11.3	9.2	11.9	13.2	12.6	11.2	10.6
2,500 – 4,499	86.4	84.5	86.8	84.3	81.0	80.5	85.3	85.6
4,500 and over	1.4	1.8	1.7	1.2	n.p.	n.p.	1.2	1.4
Not stated	—	0.1	—	—	n.p.	n.p.	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
<i>Less than 2,500</i>	<i>12.2</i>	<i>13.6</i>	<i>11.5</i>	<i>14.5</i>	<i>17.6</i>	<i>19.1</i>	<i>13.5</i>	<i>13.0</i>

(a) Care must be taken when interpreting the ACT rates as these rates include babies born in the Australian Capital Territory to Indigenous mothers who were non-ACT residents. For example, the percentage of low birthweight among babies who were born in the Australian Capital Territory to ACT-resident Indigenous mothers from 2001 to 2004 was 12.3%.

Note: Data for Tasmania were not available.

n.p. Data not published to maintain confidentiality of small numbers.

Source: AIHW NPSU National Perinatal Data Collection 2007.

In 2001–2004, the mean birthweight of live babies born to Indigenous mothers decreased with increased remoteness of the mothers' usual place of residence (Table 4.6). The mean birthweight was highest among babies born to Indigenous mothers residing in major cities

(3,188 grams), decreasing with increased remoteness to 3,169 grams in outer regional areas and 3,123 grams in very remote areas.

The proportion of liveborn babies to Indigenous mothers who had a birthweight of less than 2,500 grams was similar across all remoteness areas (between 13% and 14%).

**Table 4.6: Live births to Indigenous mothers by birthweight and remoteness area of usual residence, 2001–2004**

Birthweight (g)	Major cities	Inner regional	Outer regional	Remote	Very remote	Total <sup>(a)</sup>
Mean	3,188	3,175	3,169	3,146	3,123	3,163
			<b>Number</b>			
Less than 1,500	196	134	236	99	174	839
1,500 – 2,499	866	607	991	471	794	3,729
2,500 – 4,499	7,159	5,002	8,431	3,388	6,177	30,157
4,500 and over	126	88	123	73	91	501
Not stated	—	1	8	—	1	10
<b>Total</b>	<b>8,347</b>	<b>5,832</b>	<b>9,789</b>	<b>4,031</b>	<b>7,237</b>	<b>35,236</b>
<i>Less than 2,500</i>	<i>1,062</i>	<i>741</i>	<i>1,227</i>	<i>570</i>	<i>968</i>	<i>4,568</i>
			<b>Per cent</b>			
Less than 1,500	2.3	2.3	2.4	2.5	2.4	2.4
1,500 – 2,499	10.4	10.4	10.1	11.7	11.0	10.6
2,500 – 4,499	85.8	85.8	86.1	84.0	85.4	85.6
4,500 and over	1.5	1.5	1.3	1.8	1.3	1.4
Not stated	—	—	0.1	—	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
<i>Less than 2,500</i>	<i>12.7</i>	<i>12.7</i>	<i>12.5</i>	<i>14.1</i>	<i>13.4</i>	<i>13.0</i>

**Table 4.7: Live births to Indigenous mothers by birthweight and socioeconomic status, 2001–2004**

Birthweight (g)	Quintile of socioeconomic disadvantage <sup>(a)</sup>					Not stated	Total
	1st quintile (most disadvantaged)	2nd quintile	3rd quintile	4th quintile	5th quintile (least disadvantaged)		
Mean	3,140	3,159	3,184	3,206	3,200	3,103	3,162
	<b>Number</b>						
Less than 1,500 g	334	211	155	99	29	13	841
1,500 – 2,499 g	1,445	1,028	670	398	113	83	3,737
2,500 – 4,499 g	11,301	8,325	5,770	3,290	1,001	482	30,169
4,500 g and over	173	141	81	78	18	10	501
Not stated	6	4	—	—	—	—	10
<b>Total</b>	<b>13,259</b>	<b>9,709</b>	<b>6,676</b>	<b>3,865</b>	<b>1,161</b>	<b>588</b>	<b>35,258</b>
<i>Less than 2,500 g</i>	<i>1,779</i>	<i>1,239</i>	<i>825</i>	<i>497</i>	<i>142</i>	<i>96</i>	<i>4,578</i>
	<b>Per cent</b>						
Less than 1,500 g	2.5	2.2	2.3	2.6	2.5	2.2	2.4
1,500 – 2,499 g	10.9	10.6	10.0	10.3	9.7	14.1	10.6
2,500 – 4,499 g	85.2	85.7	86.4	85.1	86.2	82.0	85.6
4,500 g and over	1.3	1.5	1.2	2.0	1.6	1.7	1.4
Not stated	—	—	—	—	—	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
<i>Less than 2,500</i>	<i>13.4</i>	<i>12.8</i>	<i>12.4</i>	<i>12.9</i>	<i>12.2</i>	<i>16.3</i>	<i>13.0</i>

(a) Socioeconomic status quintiles are determined by postcode of usual residence based on the Australian Bureau of Statistics SEIFA .

Source: AIHW NPSU National Perinatal Data Collection 2007.

In 2001–2004 the mean birthweight of live babies born to Indigenous mothers was lowest among teenage mothers (3,094 grams) and highest in mothers aged 25–29 years (3,200 grams and 30–34 years (3,210 grams) (Table 4.8).

The proportion of liveborn low birthweight births to Indigenous mothers was highest in older mothers (i.e. those aged 35 years and over) at 15%, and lowest among mothers aged 25–29 years (12%).

**Table 4.8: Live births to Indigenous mothers by birthweight and maternal age, 2001–2004**

Birthweight	Maternal age						Total
	Less than 20	20–24	25–29	30–34	35 and over	Not stated	
Mean	3,094	3,157	3,200	3,210	3,165	3,200	3,162
	<b>Number</b>						
Less than 1,500	183	259	183	136	80	—	841
1,500 – 2,499	868	1,108	863	587	311	—	3,737
2,500 – 4,499	6,776	9,447	7,241	4,595	2,097	13	30,169
4,500 and over	49	130	130	133	59	—	501
Not stated	2	3	2	2	1	—	10
<b>Total</b>	<b>7,878</b>	<b>10,947</b>	<b>8,419</b>	<b>5,453</b>	<b>2,548</b>	<b>13</b>	<b>35,258</b>
<i>Less than 2,500</i>	<i>1,051</i>	<i>1,367</i>	<i>1,046</i>	<i>723</i>	<i>391</i>	—	<i>4,578</i>
	<b>Per cent</b>						
Less than 1,500	2.3	2.4	2.2	2.5	3.1	—	2.4
1,500 – 2,499	11.0	10.1	10.3	10.8	12.2	—	10.6
2,500 – 4,499	86.0	86.3	86.0	84.3	82.3	100.0	85.6
4,500 and over	0.6	1.2	1.5	2.4	2.3	—	1.4
Not stated	—	—	—	—	—	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
<i>Less than 2,500</i>	<i>13.3</i>	<i>12.5</i>	<i>12.4</i>	<i>13.3</i>	<i>15.3</i>	—	<i>13.0</i>

Source: AIHW NPSU National Perinatal Data Collection 2007.

For those states and territories where smoking status data were available, the mean birthweight of live babies born to Indigenous mothers in 2001–2004 was considerably lower among mothers who smoked during pregnancy compared with mothers who did not smoke (Table 4.9). The mean birthweight of live babies born to Indigenous mothers who smoked during pregnancy was 3,037 grams, 253 grams lighter than the mean birthweight of babies of Indigenous mothers who did not smoke (3,290 grams). For live births of non-Indigenous mothers who smoked the mean birthweight was 3,210 grams, compared with 3,416 grams for babies of non-Indigenous mothers who did not smoke.



The mean birthweight of live babies born to Indigenous mothers who smoked at all during pregnancy varied among the states and territories, from 2,872 grams in the Australian Capital Territory to 3,061 grams in New South Wales.

**Table 4.9: Mean birthweight (g) of live births to Indigenous mothers by tobacco smoking status during pregnancy and state and territory, 2001–2004**

Smoking status	NSW	WA	SA <sup>(a)</sup>	ACT	NT <sup>(b)</sup>	Total
Smoked	3,061	3,015	2,955	2,872	3,057	3,037
Did not smoke	3,371	3,230	3,290	3,261	3,243	3,290
<b>Total<sup>(c)</sup></b>	<b>3,193</b>	<b>3,124</b>	<b>3,064</b>	<b>3,085</b>	<b>3,117</b>	<b>3,144</b>

(a) For South Australia, 'Smoked' includes women who quit before the first antenatal visit.

(b) For the Northern Territory, smoking status was recorded at the first antenatal visit.

(c) Includes records where smoking status was not stated.

Note: Data on tobacco smoking during pregnancy were not available for Victoria, Queensland or Tasmania.

Source: AIHW NPSU National Perinatal Data Collection 2007.

The proportion of low birthweight among liveborn babies of Indigenous mothers who smoked was 16%, compared with 10% among babies of Indigenous mothers who did not smoke. For non-Indigenous mothers, these figures were 10% and 5%, respectively.

## Apgar scores

Apgar scores are clinical indicators of the baby's condition shortly after birth, based on assessment of the heart rate, breathing, colour, muscle tone and reflex irritability. Between 0 and 2 points are given for each of these five characteristics, and the total score is between 0 and 10. An Apgar score of less than 7 at 5 minutes after birth is considered to be an indicator of compromise for the baby.

In 2001–2004, around 3% of liveborn babies to Indigenous mothers had a low Apgar score (between 0 and 6) at 5 minutes (Table 4.10). This proportion was greater than the proportion in non-Indigenous mothers (1.3%).

Among the states and territories, the distribution of low Apgar scores at 5 minutes for live babies born to Indigenous mothers ranged from 2% in New South Wales, Victoria, Queensland and Western Australia to 4% in the Northern Territory.

**Table 4.10: Live births to Indigenous mothers by Apgar score at 5 minutes and state and territory, 2001–2004**

Apgar score	NSW	Vic	Qld	WA	SA	ACT	NT	Total
	<b>Number</b>							
0–6	196	34	229	139	44	7	216	865
7–10	8,516	1,587	10,792	5,998	1,732	269	5,370	34,264

# Hospital births

## Length of stay

The majority of babies are discharged from hospital at the same time as their mothers; however, some babies experience morbidity and require hospitalisation. A baby's gestation and birthweight are two factors that influence the duration of hospitalisation. Twins and higher order multiple births usually have longer stays in hospital than singleton babies.

In 2001–2004, the median length of stay for babies born to Indigenous mothers in hospital who were discharged home was 3 days (Table 4.12). The majority of babies born to Indigenous mothers remained in their hospital of birth for less than 6 days (84%), and nearly two-thirds stayed in hospital for less than 4 days (61%). Length of stay in hospital for babies born to Indigenous mothers was generally shorter when compared with babies born to non-Indigenous mothers. Only 45% of babies born in hospital to non-Indigenous mothers had a length of stay of less than 4 days.

There was some variation among the states and territories in the length of stay for babies born to Indigenous mothers who were discharged from hospital in 2001–2004. Over one-fifth of babies born to Indigenous mothers in hospital in the Northern Territory had a length of stay of 7 days or more (21%) compared with only 10% in Victoria. The proportion of babies born to Indigenous mothers who had a length of stay of less than 4 days ranged from 43% in the Northern Territory to 71% in Victoria.

**Table 4.12: Live births to Indigenous mothers in hospital<sup>(a)</sup> by length of stay and state and territory, 2001–2004**

Length of stay	NSW	Vic	Qld	WA	SA	ACT	NT	Total
Median length of stay	3.0	3.0	2.0	3.0	3.0	3.0	4.0	3.0
	<b>Number</b>							
Less than 1 day	228	39	441	113	45	8	43	917
1 day	1,222	189	2,385	753	166	31	279	5,025
2 days	1,763	343	2,552	1,175	330	44	736	6,943
3 days	1,640	357	1,800	1,087	305	37	890	6,116
4 days	1,077	216	1,105	901	238	34	712	4,283
5 days	695	136	594	578	137	31	583	2,754
6 days	327	51	320	283	77	11	318	1,387
7–13 days	492	82	536	468	127	17	610	2,332
14–27 days	171	44	277	123	60	6	201	882
28 or more days	86	23	154	65	39	9	145	521
Not stated	7	—	—	—	—	—	—	7
<b>Total</b>	<b>7,708</b>	<b>1,480</b>	<b>10,164</b>	<b>5,546</b>	<b>1,524</b>	<b>228</b>	<b>4,517</b>	<b>31,167</b>
	<b>Per cent</b>							
Less than 1 day	3.0	2.6	4.3	2.0	3.0	3.5	1.0	2.9
1 day	15.9	12.8	23.5	13.6	10.9	13.6	6.2	16.1
2 days	22.9	23.2	25.1	21.2	21.7	19.3	16.3	22.3
3 days	21.3	24.1	17.7	19.6	20.0	16.2	19.7	19.6
4 days	14.0	14.6	10.9	16.2	15.6	14.9	15.8	13.7
5 days	9.0	9.2	5.8	10.4	9.0	13.6	12.9	8.8
6 days	4.2	3.4	3.1	5.1	5.1	4.8	7.0	4.5
7–13 days	6.4	5.5	5.3	8.4	8.3	7.5	13.5	7.5
14–27 days	2.2	3.0	2.7	2.2	3.9	2.6	4.4	2.8
28 or more days	1.1	1.6	1.5	1.2	2.6	3.9	3.2	1.7
Not stated	0.1	—	—	—	—	—	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Only babies who were discharged home are included.

Note: Data for Tasmania were not available.

Source: AIHW NPSU National Perinatal Data Collection 2007.

Compared with other remoteness areas, the length of stay for babies born to Indigenous mothers who were discharged from hospital was considerably higher where the mothers' usual place of residence was in very remote areas (Table 4.13). In 2001–2004, babies born to Indigenous mothers whose usual place of residence was in very remote areas accounted for almost one-fifth of babies discharged from hospital with a length of stay of 7 days or more (19%). This compared with 12% in remote areas, 12% in major cities, 9% in outer regional areas and 8% in inner regional areas. The proportion of babies born to Indigenous mothers who were discharged from hospital after a length of stay of less than 4 days ranged from 81% for mothers residing in very remote areas to 92% for mothers residing in inner regional areas.

**Table 4.13: Live births to Indigenous mothers in hospital<sup>(a)</sup> by length of stay and remoteness area of usual residence, 2001–2004**

Length of Stay	Major cities	Inner regional	Outer regional	Remote	Very remote	Total <sup>(b)</sup>
Median length of stay	3.0	3.0	3.0	3.0	4.0	3.0
<b>Number</b>						
Less than 1 day	274	172	325	77	68	916
1 day	1,271	982	1,706	547	519	5,025
2 days	1,705	1,286	2,124	802	1,023	6,940
3 days	1,423	1,133	1,669	702	1,187	6,114
4 days	1,061	710	1,050	471	990	4,282
5 days	675	391	651	282	754	2,753
6 days	308	178	306	145	450	1,387
7–13 days	503	253	480	265	827	2,328
14–20 days	148	102	128	59	166	603
21–27 days	100	43	52	29	53	277
28 or more days	166	48	132	62	111	519
Not stated	2	1	2	1	1	7
<b>Total</b>	<b>7,636</b>	<b>5,299</b>	<b>8,625</b>	<b>3,442</b>	<b>6,149</b>	<b>31,151</b>
<b>Per cent</b>						
Less than 1 day	3.6	3.2	3.8	2.2	1.1	2.9
1 day	16.6	18.5	19.8	15.9	8.4	16.1
2 days	22.3	24.3	24.6	23.3	16.6	22.3
3 days	18.6	21.4	19.4	20.4	19.3	19.6
4 days	13.9	13.4	12.2	13.7	16.1	13.7
5 days	8.8	7.4	7.5	8.2	12.3	8.8
6 days	4.0	3.4	3.5	4.2	7.3	4.5
7–13 days	6.6	4.8	5.6	7.7	13.4	7.5
14–20 days	1.9	1.9	1.5	1.7	2.7	1.9
21–27 days	1.3	0.8	0.6	0.8	0.9	0.9

-1812(

second most disadvantaged quintile (second quintile) to 13% in the second least disadvantaged quintile (fourth quintile).

**Table 4.14: Live births to Indigenous mothers in hospitals<sup>(a)</sup>, by length of stay and socioeconomic status, 2001–2004**

Length of stay	Quintile of socioeconomic disadvantage <sup>(b)</sup>					Not stated	Total
	1st quintile (most disadvantaged)	2nd quintile	3rd quintile	4th quintile	5th quintile (least disadvantaged)		
Median length of stay	3.0	3.0	3.0	3.0	4.0	3.0	3.0
	<b>Number</b>						
Less than 1 day	332	265	192	97	27	4	917
1 day	1,701	1,418	1,193	556	116	41	5,025
2 days	2,367	2,034	1,466	755	210	111	6,943
3 days	2,287	1,800	1,116	656	172	85	6,116
4 days	1,541	1,163	806	528	169	76	4,283
5 days	1,077	701	487	325	123	41	2,754
6 days	589	346	228	128	70	26	1,387
7–13 days	936	554	410	263	113	56	2,332
14–20 days	224	151	122	71	26	10	604
21–27 days	97	58	60	38	14	11	278
28 or more days	176	129	99	83	23	11	521
Not stated	5	2	—	—	—	—	7
<b>Total</b>	<b>11,332</b>	<b>8,621</b>	<b>6,179</b>	<b>3,500</b>	<b>1,063</b>	<b>472</b>	<b>31,167</b>
	<b>Per cent</b>						
Less than 1 day	2.9	3.1	3.1	2.8	2.5	0.8	2.9
1 day	15.0	16.4	19.3	15.9	10.9	8.7	16.1
2 days	20.9	23.6	23.7	21.6	19.8	23.5	22.3
3 days	20.2	20.9	18.1	18.7	16.2	18.0	19.6
4 days	13.6	13.5	13.0	15.1	15.9	16.1	13.7
5 days	9.5	8.1	7.9	9.3	11.6	8.7	8.8
6 days	5.2	4.0	3.7	3.7	6.6	5.5	4.5
7–13 days	8.3	6.4	6.6	7.5	10.6	11.9	7.5
14–20 days	2.0	1.8	2.0	2.0	2.4	2.1	1.9
21–27 days	0.9	0.7	1.0	1.1	1.3	2.3	0.9
28 or more days	1.6	1.5	1.6	2.4	2.2	2.3	1.7
Not stated	—	—	—	—	—	—	—
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Only babies who were discharged home are included.

# Perinatal mortality

## Definitions

There are different definitions in Australia for reporting and registering perinatal deaths. The NHDD specifies a definition of perinatal deaths to include all fetal and neonatal deaths of at least 400 grams birthweight or at least 20 weeks gestation (NHDC 2003). This definition is used in the NPDC.

Figure 4.1 shows the definitions of periods of perinatal and infant deaths used by the NPSU. Neonatal deaths are those occurring in live births up to 28 completed days after birth.

Labour		Birth	7 days	28 days	1 year
<i>At least 20 weeks or 400 grams</i>		<i>0-7 days</i>	<i>7-28 days</i>	<i>28 days-1 year</i>	
Antepartum fetal deaths	Intrapartum fetal deaths	Early neonatal deaths	Late neonatal deaths	Postneonatal deaths	
<b>Fetal deaths</b>		<b>Neonatal deaths</b>			
<b>Perinatal deaths</b>					
		<b>Infant deaths</b>			

**Figure 4.1: Perinatal and infant death periods**

**Table 4.15: Fetal, neonatal and perinatal deaths of babies born to Indigenous mothers by state and territory, 2001–2004**

Deaths	NSW	Vic	Qld	WA	SA	ACT <sup>(a)</sup>	NT	Total <sup>(b)</sup>
	<b>Number</b>							
Fetal deaths	91	28	121	84	30	7	63	424
Neonatal deaths	32	15	88	46	15	5	n.a.	201
<b>Perinatal deaths</b>	<b>123</b>	<b>43</b>	<b>209</b>	<b>130</b>	<b>45</b>	<b>12</b>	<b>n.a.</b>	<b>625</b>
	<b>Rate per 1,000 births<sup>(c)</sup></b>							
Fetal deaths	10.3	16.9	10.8	13.4	16.5	24.6	11.1	11.9
Neonatal deaths	3.7	9.2	8.0	7.5	8.4	18.1	n.a.	5.7
<b>Perinatal deaths</b>	<b>13.9</b>	<b>26.0</b>	<b>18.7</b>	<b>20.8</b>	<b>24.8</b>	<b>42.3</b>	<b>n.a.</b>	<b>17.5</b>

(a) Care must be taken when interpreting these rates as the small number of perinatal deaths in the Australian Capital Territory will result in large fluctuations over time. ACT rates include babies born in the Australian Capital Territory to Indigenous mothers who were non-ACT residents. There were four fetal deaths (19.2 per 1,000 births), no neonatal deaths and a perinatal death rate of 19.2 per 1,000 births for babies born to ACT-resident Indigenous women in 2001–2004.

(b) Totals for neonatal and perinatal deaths exclude neonatal deaths in the Northern Territory.

(c) Fetal and perinatal death rates were calculated using all births (live births and still births). Neonatal death rates were calculated using all live births.

n.a. Data on neonatal deaths were not available for the Northern Territory for 2001–2003, therefore neonatal deaths in the Northern Territory have not been included.

Source: AIHW NPSU National Perinatal Data Collection 2007.



## **5 Assessment of Indigenous status data quality, 1991 to 2004**

### **Summary**

An assessment of the quality of Indigenous status information in perinatal data in each state and territory was undertaken. This involved a survey which was sent to the midwifery managers across Australia to determine how many hospitals in each jurisdiction obtain Indigenous status information of mothers giving birth from admission records and how many collect this information independently. The assessment also involved analysis of the variability in the number and proportion of mothers recorded as Indigenous in the perinatal data collection over time and across jurisdictions for the period 1991–2004 (Table 5.3).

The outcomes of this assessment showed that Indigenous status data from New South Wales, Victoria, Queensland, Western Australia, South Australia and the Northern Territory are suitable for trends analysis. Perinatal data from Tasmania, although improving, were deemed to be of insufficient quality. Although the most recent data in the Australian Capital

resumed after various problems were resolved including the addition of a specific question regarding the Indigenous status of the babies' fathers. The percentage of obstetric patients who reported being asked their Indigenous status improved markedly after the resumption of the study. The study also found that there were a number of inconsistencies in the group of women who identified themselves or their babies as Indigenous with the medical records (Middleton et al. 2003).

A similar study was conducted by the University of New South Wales between May and July 1999 (Jackson Pulver et al. 2003). The study attempted to determine the accuracy of routine identification of Aboriginal and Torres Strait Islander women admitted to King George V hospital in Sydney to give birth. Hospital records were compared with self-disclosure of Aboriginal and Torres Strait Islander status to a female Aboriginal health professional.

Of the 536 women surveyed, 29 (5%) self-disclosed as being Indigenous. Only a third of the patients were identified as Aboriginal and Torres Strait Islander in hospital records. The Indigenous women referred by another organisation were significantly more likely to be correctly identified than those who self-referred to the hospital (Jackson Pulver et al. 2003).

In New South Wales, the quality of Indigenous status in perinatal data has also been assessed through linking birth records reported to the NSW Midwives Data Collection (MDC) and birth registrations reported to the NSW Registry of Births, Deaths and Marriages in the period 1994–1998. Records using the two databases were matched using probabilistic linkage





information checked by midwives. The other six jurisdictions generally have more sources to obtain Indigenous status information, or more frequently have the information checked by midwives.

**Table 5.1: How the Indigenous status of the mother is obtained, by state and territory, 2007**

	NSW	Vic	Qld	WA	SA	Tas	ACT <sup>(a)</sup>	NT	Total
<b>Number</b>									
Always from hospital database/admission form	21	25	8	5	13	2	2	1	77
Always independent from hospital database	32	11	22	6	6	1	—	2	80
Collected by midwife only if information is not recorded in hospital database	6	6	8	9	5	—	—	—	34
Other <sup>(b)</sup>	5	1	5	3	2	—	—	—	16
<b>Total</b>	<b>64</b>	<b>43</b>	<b>43</b>	<b>23</b>	<b>26</b>	<b>3</b>	<b>2</b>	<b>3</b>	<b>207</b>
<b>Per cent</b>									
Always from hospital database/admission form	32.8	58.1	18.6	21.7	50.0	66.7	100.0	33.3	37.2
Always independent from hospital database	50.0	25.6	51.2	26.1	23.1	33.3	0.0	66.7	38.6
Collected by midwife only if information is not recorded in hospital database	9.4	14.0	18.6	39.1	19.2	0.0	0.0	0.0	16.4
Other <sup>(b)</sup>	7.8	2.3	11.6	13.0	7.7	0.0	0.0	0.0	7.7
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Survey was administered by telephone in the Australian Capital Territory. Results should be interpreted with caution as these are based on only two respondents.

(b) Includes: collected from both the hospital database and by midwives (8 respondents); collected on booking in form; Indigenous status of both parents is recorded and checked at all of the above steps (inc10(cordeoTc -0.0045 Tw 3.0009 0 Tdene baapitclid onic, atio(i)(s aatu)s-5(io)0(n)to bi7.2)TJ0.0003 To



from 1993 to 2004. Table 5.4 shows the variability in the numbers and proportions of Indigenous non-residents from year to year, compared with little variability among Indigenous ACT residents who give birth in the Australian Capital Territory.

For the other jurisdictions, the number of mothers identified as Indigenous followed smooth trends over time, either increasing at a relatively steady rate (New South Wales, Queensland, and the Northern Territory), or remaining at a relatively stable level (Victoria, Western Australia and South Australia).

Jurisdictions generally started to record mothers with a 'not stated' Indigenous status in or after 1999. In Western Australia and South Australia, no mothers with a 'not stated' Indigenous status were recorded in the period 1991–2004.

In Western Australia, data for the mother's Indigenous status were drawn from two sources—(1) from the mother's record in the Hospital Morbidity Data Collection, and (2) if these data were missing on the inpatient record or the mother did not give birth in hospital, from the form completed by the midwife (Notification of Case Attended form). This form does not allow for a 'not stated' option and a valid response category must be recorded.





**Table 5.3 (continued): Women who gave birth by Indigenous status<sup>(a)</sup> and state and territory, 1991 to 2004**

	Indigenous status <sup>(a)</sup>	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
<b>SA</b>															
	Indigenous	397	435	396	388	382	348	393	412	445	447	398	443	468	484
	Non-Indigenous	19,071	19,433	19,283	19,131	18,928	18,436	18,001	18,008	17,788	17,131	17,029	16,978	17,049	16,744
	Not stated	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	<b>Total</b>	<b>19,468</b>	<b>19,868</b>	<b>19,679</b>	<b>19,519</b>	<b>19,310</b>	<b>18,784</b>	<b>18,394</b>	<b>18,420</b>	<b>18,233</b>	<b>17,578</b>	<b>17,427</b>	<b>17,421</b>	<b>17,517</b>	<b>17,228</b>
<b>ACT</b>															
	Indigenous	20	11	48	44	63	80	58	63	59	53	52	72	80	73
	Non-Indigenous	4,459	4,632	4,664	4,686	4,767	4,621	4,650	4,582	4,522	4,625	4,353	4,610	4,683	4,711
	Not stated	—	—	—	—	—	—	—	—	5	6	9	26	21	15
	<b>Total</b>	<b>4,479</b>	<b>4,643</b>	<b>4,712</b>	<b>4,730</b>	<b>4,830</b>	<b>4,701</b>	<b>4,708</b>	<b>4,645</b>	<b>4,586</b>	<b>4,684</b>	<b>4,414</b>	<b>4,708</b>	<b>4,784</b>	<b>4,799</b>
<b>NT</b>															
	Indigenous	1,209	1,243	1,221	1,220	1,244	1,210	1,197	1,248	1,295	1,348	1,485	1,409	1,396	1,332
	Non-Indigenous	2,254	2,343	2,286	2,260	2,363	2,212	2,328	2,240	2,252	2,273	2,228	2,233	2,216	2,096
	Not stated	—	—	—	—	—	—	—	—	—	8	16	32	6	9
	<b>Total</b>	<b>3,463</b>	<b>3,586</b>	<b>3,507</b>	<b>3,480</b>	<b>3,607</b>	<b>3,422</b>	<b>3,525</b>	<b>3,488</b>	<b>3,547</b>	<b>3,629</b>	<b>3,729</b>	<b>3,674</b>	<b>3,618</b>	<b>3,437</b>
<b>Australia<sup>(b)</sup></b>															
	<b>Indigenous</b>	<b>7,025</b>	<b>7,253</b>	<b>7,249</b>	<b>7,390</b>	<b>7,774</b>	<b>7,815</b>	<b>7,894</b>	<b>8,446</b>	<b>8,697</b>	<b>8,628</b>	<b>8,681</b>	<b>8,822</b>	<b>8,857</b>	<b>8,904</b>
	<b>Non-Indigenous</b>	<b>239,256</b>	<b>244,974</b>	<b>242,830</b>	<b>243,485</b>	<b>241,921</b>	<b>238,916</b>	<b>239,098</b>	<b>236,991</b>	<b>238,622</b>	<b>238,564</b>	<b>235,699</b>	<b>236,194</b>	<b>238,214</b>	<b>238,466</b>
	<b>Not stated</b>	<b>—</b>	<b>—</b>	<b>—</b>	<b>—</b>	<b>—</b>	<b>—</b>	<b>—</b>	<b>—</b>	<b>21</b>	<b>72</b>	<b>79</b>	<b>117</b>	<b>76</b>	<b>88</b>
	<b>Total</b>	<b>246,281</b>	<b>252,227</b>	<b>250,079</b>	<b>250,875</b>	<b>249,695</b>	<b>246,731</b>	<b>246,992</b>	<b>245,437</b>	<b>247,340</b>	<b>247,264</b>	<b>244,459</b>	<b>245,133</b>	<b>247,147</b>	<b>247,458</b>

(a) The categories used for recording Indigenous status are based on the National Health Data Dictionary value domains. From 1999 onwards the recording categories used were

**Table 5.4: Resident and non-resident women who gave birth in the Australian Capital Territory by Indigenous status, 1993 to 2004**

	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
<b>ACT residents</b>						<b>Number</b>						
Indigenous	53	50	59	67	51	47	50	38	40	50	58	54
Non-Indigenous	4,097	4,192	4,306	4,167	4,053	4,003	3,992	4,060	3,762	3,907	3,979	3,951
<b>Total</b>	<b>4,150</b>	<b>4,242</b>	<b>4,365</b>	<b>4,234</b>	<b>4,104</b>	<b>4,050</b>	<b>4,042</b>	<b>4,098</b>	<b>3,802</b>	<b>3,957</b>	<b>4,037</b>	<b>4,005</b>
						<b>Per cent</b>						
Indigenous	1.3	1.2	1.4	1.6	1.2	1.2	1.2	0.9	1.1	1.3	1.4	1.3
Non-Indigenous	98.7	98.8	98.6	98.4	98.8	98.8	98.8	99.1	98.9	98.7	98.6	98.7
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
<b>Non-ACT residents</b>						<b>Number</b>						
Indigenous	10	6	5	14	7	16	9	15	12	22	22	19
Non-Indigenous	439	495	509	510	597	579	530	565	591	703	704	760
<b>Total</b>	<b>449</b>	<b>501</b>	<b>514</b>	<b>524</b>	<b>604</b>	<b>595</b>	<b>539</b>	<b>580</b>	<b>603</b>	<b>725</b>	<b>726</b>	<b>779</b>
						<b>Per cent</b>						
Indigenous	2.2	1.2	1.0	2.7	1.2	2.7	1.7	2.6	2.0	3.0	3.0	2.4
Non-Indigenous	97.8	98.8	99.0	97.3	98.8	97.3	98.3	97.4	98.0	97.0	97.0	97.6
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

*Notes*


1. Excludes records where Indigenous status or state of usual residence were 'Not stated.'
2. Data before 1993 not available in the required format.

Source: ACT Maternal Perinatal Data Collection.

## 2. Proportion of Indigenous mothers

The proportion of women who gave birth and identified as Indigenous between 1991 and 2004 are presented by state and territory in Figures 5.1a and 5.1b.



Similar to the raw data provided in Table 5.3, the proportions of Indigenous mothers showed a variable pattern in the Australian Capital Territory over the period 1991–2004. There was less variation when only ACT-resident Indigenous mothers were considered (Table 5.4). In the other states and territories, the proportion of Indigenous mothers remained relatively stable over this period.



*Note: The scale used for the Northern Territory is different to that used for 4.658*

### **3. Variability of the number of Indigenous mothers based on coefficient of variation**

Over the period 1991–2004, the number of mothers identified as Indigenous was most variable in the Australian Capital Territory compared with other states and territories, although this has become more stable in later years. In Figure 5.2b, the coefficient of variation in the number of Indigenous mother



Note: Coefficient of variation was calculated based on the number of Indigenous mothers for every five-year period from 1991-1995 to 2000-2004.

Source: AIHW NPSU National Perinatal Data Collection 2007.

**Figure 5.2b: Coefficient of variation for the number of Indigenous mothers, by state and territory, 1991-1995 to 2000-2004**







## 6 Trends

### Summary

Although Indigenous mothers remain considerably younger than non-Indigenous mothers, there was a trend over the 1991–2004 period of

Long-term trends for each of the variables presented in this chapter have been analysed by modelling trends throughout the period 1991–2004. A linear regression model has been used for the analysis of trends which takes into account all observations throughout the period and fits a line of best fit to the data. The average annual change occurring from year to year over the period has been calculated based on the fitted trend. When the 99% confidence intervals for the trend (p-value of less than 0.01) both lie above or below zero, the trend is characterised in subsequent text as a ‘significant’ increase or a ‘significant’ decline.

The trends analyses presented in this chapter on the proportion of mothers who have given birth by caesarean section have used age-standardised data. Age-standardisation allows for comparisons between populations which have different age structures, as is the case with Indigenous and non-Indigenous mothers. All age-standardised rates and rate ratios have been calculated using the direct standardisation method and ‘all women who gave birth in Australia in 2001’ has been used as the standard population. The Appendix contains further detail on these methods.

As discussed in Chapter 5, fluctuations in the proportion of Indigenous mothers over time may partly reflect changing levels of coverage of Indigenous mothers in the perinatal data. Therefore, caution should be used when assessing trends over time and when making comparisons with non-Indigenous mothers.

## Maternal age

Table 6.1 presents the rate of Indigenous and non-Indigenous women who gave birth per 100 women in the reproductive population and the rate ratios of Indigenous and non-Indigenous mothers over the period 1991–2004.

The proportion of Indigenous women in the population who gave birth during the 1991–2004 period remained stable and the proportion of non-Indigenous women who gave birth over this period decreased significantly (Table 6.1). The fitted trend indicated an average yearly decrease of four non-Indigenous mothers per 10,000 population.

There was a trend towards increasing proportions of older mothers and decreasing proportions of younger mothers in both Indigenous and non-Indigenous mothers during the 1991–2004 period. Among Indigenous women, there was a significant decrease in the proportion of teenage mothers and a significant increase in the proportion of mothers aged 25–29 years, 30–34 years and 35 years and over.

Among non-Indigenous women, there was a significant decrease in teenage mothers and those aged 20–24 years and 25–29 years. There was a significant increase in the proportion of non-Indigenous mothers aged 30–34 years and 35 years and over.

The rate ratios between Indigenous and non-Indigenous mothers increased significantly for those aged less than 20 years, 20–24 years, 25–29 years and 30–34 years over the 1991–2004 period. The proportion of teenage mothers decreased among both Indigenous and non-Indigenous women; however, the decrease was greater for Indigenous teenage mothers (an average annual decrease of 0.13%) compared with non-Indigenous mothers (an average annual decrease of 0.04%). The proportion of Indigenous mothers aged 20–24 years did not change significantly over time but non-Indigenous mothers in this age group decreased significantly by a yearly average of 0.16%. The proportion of Indigenous mothers aged 25–29 years increased significantly (by an annual average of 14 mothers per 10,000

population) whereas the proportion of non-Indigenous mothers in this age group decreased (by an annual average of 24 mothers per 10,000 population).

The proportions of both Indigenous and non-Indigenous mothers aged 30–34 years increased significantly over the period; however, the increase was greater in Indigenous mothers.

**Table 6.1: Rates (proportions)<sup>(a)</sup> and rate ratios of Indigenous and non-Indigenous mothers by maternal age, 1991 to 2004**

Maternal age	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	Annual change <sup>(b)</sup>
<b>Indigenous rate per 100 women <sup>(a)</sup>(%)</b>															
Less than 20	10.0	9.9	9.4	10.2	9.9	9.5	9.8	9.3	9.5	9.4	8.9	8.7	8.5	8.1	-0.13
20–24	14.3	14.0	13.6	13.8	14.3	14.2	14.2	14.6	15.4	14.5	14.9	14.8	14.3	13.9	0.05
25–29	9.7	10.3	10.5	10.1	11.6	11.2	11.1	12.0	12.2	11.6	11.7	11.6	11.7	11.3	0.14*
30–34	5.6	6.0	6.1	5.8	5.8	6.3	6.3	7.5	7.2	7.4	7.5	7.2	7.2	7.5	0.16*
35 and over	1.4	1.6	1.5	1.5	1.6	1.7	1.7	2.0	2.0	2.2	1.9	2.2	2.1	2.3	0.07*
<b>Total</b>	<b>8.2</b>	<b>8.3</b>	<b>8.1</b>	<b>8.1</b>	<b>8.4</b>	<b>8.3</b>	<b>8.2</b>	<b>8.6</b>	<b>8.7</b>	<b>8.5</b>	<b>8.3</b>	<b>8.3</b>	<b>8.1</b>	<b>8.0</b>	<b>0.00</b>
<b>Non-Indigenous rate per 100 women <sup>(a)</sup>(%)</b>															
Less than 20	2.0	2.0	1.9	2.0	1.9	1.9	1.9	1.8	1.8	1.7	1.6	1.6	1.5	1.5	-0.04*
20–24	7.2	7.2	7.0	6.9	6.6	6.4	6.2	6.1	6.1	5.9	5.8	5.5	5.3	5.2	-0.16*
25–29	12.9	13.0	12.8	12.6	12.2	11.7	11.4	11.2	11.1	10.9	10.6	10.4	10.3	10.1	-0.24*
30–34	9.9	10.3	10.4	10.4	10.6	10.5	10.7	10.7	10.9	11.0	10.9	11.1	11.3	11.4	0.10*
35 and over	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.8	2.9	2.9	3.0	3.2	3.3	0.09*
<b>Total</b>	<b>6.2</b>	<b>6.3</b>	<b>6.2</b>	<b>6.2</b>	<b>6.2</b>	<b>6.0</b>	<b>6.0</b>	<b>6.0</b>	<b>6.0</b>	<b>5.9</b>	<b>5.8</b>	<b>5.8</b>	<b>5.8</b>	<b>5.8</b>	<b>-0.04*</b>
<b>Rate ratio</b>															
Less than 20	5.0	5.0	4.9	5.2	5.1	5.0	5.2	5.2	5.4	5.6	5.5	5.5	5.8	5.5	0.06*
20–24	2.0	1.9	1.9	2.0	2.2	2.2	2.3	2.4	2.5	2.5	2.6	2.7	2.7	2.7	0.07*
25–29	0.7	0.8	0.8	0.8	1.0	1.0	1.0	1.1	1.1	1.1	1.1	1.1	1.1	1.1	0.03*
30–34	0.6	0.6	0.6	0.6	0.5	0.6	0.6	0.7	0.7	0.7	0.7	0.6	0.6	0.7	0.01*
35 and over	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.00
<b>Total</b>	<b>1.3</b>	<b>1.3</b>	<b>1.3</b>	<b>1.3</b>	<b>1.4</b>	<b>1.4</b>	<b>1.4</b>	<b>1.4</b>	<b>1.5</b>	<b>1.4</b>	<b>1.4</b>	<b>1.4</b>	<b>1.4</b>	<b>1.4</b>	<b>0.01*</b>

(a) Based on women aged 15–44 years in the population.

(b) Average annual change in number and proportion of mothers determined using linear regression analysis.

## **Age of first-time mothers**


**Table 6.2: Rates (proportions)<sup>(a)</sup> and rate ratios of Indigenous and non-Indigenous first-time mothers by maternal age, 1991 to 2004**

Maternal age	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	Annual change <sup>(b)</sup>
	<b>Indigenous rate per 100 women who gave birth (%)</b>														
Less than 20	67.8	68.6	67.4	68.8	69.3	69.4	70.2	69.1	72.8	72.8	72.2	69.7	71.6	72.4	0.36*
20–24	25.6	25.4	24.7	25.9	25.7	26.2	28.1	25.8	27.2	24.9	26.8	27.6	28.8	29.6	0.26*
25–29	12.3	11.4	12.8	12.7	12.2	12.4	13.0	13.9	13.7	12.9	12.6	11.5	13.3	12.9	0.05
30–34	8.0	9.1	7.5	9.9	8.7	8.3	9.6	10.8	11.0	9.1	9.6	10.5	9.8	10.7	0.17*
35 and over															

## Caesarean sections

The caesarean section age-standardised rate has increased significantly over the 1991–2004 period among both Indigenous and non-Indigenous mothers (Figure 6.1). Rates were directly standardised to the 2001 population of women who gave birth in order to control for the effect of maternal age on the prevalence of caesarean sections.

The Indigenous age-standardised caesarean section rate increased from 20% in 1991 to 26% in 2004, an average annual increase of 0.54% per annum (Table 6.3). The caesarean section rate among non-Indigenous women increased at a greater rate over this same period. In 1991, 19% of non-Indigenous mothers had a caesarean section, increasing to 29% in 2004. This was an average annual increase of 0.8% per annum.



**Table 6.3: Age-standardised rate (proportion)<sup>(a)</sup> and rate ratios of Indigenous and non-Indigenous women who gave birth by caesarean section, 1991 to 2004**

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	Annual change <sup>(b)</sup>
<b>Age-standardised rate per 100 women who gave birth (%)</b>															
Indigenous	19.8	19.4	20.2	20.2	21.1	19.8	20.9	21.5	21.2	22.8	24.4	24.6	26.5	26.4	0.54*
Non-Indigenous	18.8	19.1	19.7	20.0	19.9	19.9	20.6	21.4	22.2	23.5	25.5	27.1	28.3	29.0	0.80*
<b>Rate ratio</b>	<b>1.1</b>	<b>1.0</b>	<b>1.0</b>	<b>1.0</b>	<b>1.1</b>	<b>1.0</b>	<b>1.0</b>	<b>1.0</b>	<b>1.0</b>	<b>1.0</b>	<b>1.0</b>	<b>0.9</b>	<b>0.9</b>	<b>0.9</b>	<b>-0.01*</b>

(a) Standardised to the population of women who gave birth in 2001.

(b) Average annual change in number and proportion of mothers determined using linear regression analysis.

\* Represents results with a statistically significant increase or decrease over the 1991–2004 period using 99% confidence intervals.

Note: Excludes data from Tasmania and the Australian Capital Territory.


Source: AIHW NPSU National Perinatal Data Collection 2007.

## Birthweight

Figure 6.2 shows the trend in low birthweight of singleton liveborn babies over the period 1991–2004 by Indigenous status. Over the 14-year period, the proportion of low birthweight babies (i.e. less than 2,500 grams) increased significantly for babies born to both Indigenous and non-Indigenous mothers.

The proportion of low birthweight babies born to non-Indigenous mothers increased slightly from 4.5% in 1991 to 4.6% in 2004; however, this increase was significant, with an average yearly increase of 0.01 per 100 live births (Table 6.4).

The proportion of low birthweight babies born to Indigenous mothers increased at a greater rate and remained more than double the proportion seen among babies born to non-Indigenous mothers over the 1991–2004 period. In 1991, 11% of babies born to Indigenous mothers were of low birthweight. This increased to 12% in 2004, an average yearly increase of 0.12 per 100 liveborn babies.





**Table 6.4: Rates (proportion) and rate ratios of low birthweight babies<sup>(a)</sup> born to Indigenous and non-Indigenous mothers, 1991 to 2004**

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	Annual change <sup>(b)</sup>
<b>Age-standardised rate per 100 births (%)</b>															
Indigenous	11.1	10.8	9.7	11.0	10.2	10.7	11.3	10.0	10.9	11.7	11.5	11.8	11.9	12.1	0.12*
Non-Indigenous	4.5	4.4	4.5	4.4	4.5	4.5	4.5	4.6	4.6	4.5	4.5	4.6	4.5	4.6	0.01*
<b>Rate ratio</b>	<b>2.5</b>	<b>2.4</b>	<b>2.2</b>	<b>2.5</b>	<b>2.3</b>	<b>2.4</b>	<b>2.5</b>	<b>2.2</b>	<b>2.4</b>	<b>2.6</b>	<b>2.6</b>	<b>2.6</b>	<b>2.6</b>	<b>2.6</b>	<b>0.02*</b>

(a) Includes liveborn singleton babies only.

(b) Average annual change in number and proportion of mothers determined using linear regression analysis.

\* Represents results with a statistically significant increase or decrease over the 1991–2004 period using 99% confidence intervals.

Note: Excludes data from Tasmania and the Australian Capital Territory.


Source: AIHW NPSU National Perinatal Data Collection 2007.

## Gestational age

Over the 14-year period 1991–2004, the proportion of liveborn singleton preterm babies (i.e. less than 37 weeks gestation) did not change for babies born to either Indigenous mothers or non-Indigenous mothers (Figure 6.3). The fitted trend implies an average annual increase of 0.05% for babies born to Indigenous mothers and 0.02% for babies born to non-Indigenous mothers, but neither increase was statistically significant (Table 6.5).

The proportion of preterm babies born to Indigenous mothers was consistently more than double the proportion of preterm babies born to non-Indigenous mothers over time. The proportion of preterm babies born to Indigenous mothers varied from 10% to 12% over the 1991–2004 period. The proportion of preterm babies born to non-Indigenous mothers remained between 5% and 6% over the 1991–2004 period (Table 6.5).

There was no change over time in the rate ratio between the proportion of preterm babies born to Indigenous and non-Indigenous mothers (Table 6.5)




**Table 6.5: Rates (proportion) and rate ratios of preterm babies<sup>(a)</sup>**

## Fetal deaths

Figure 6.4 shows the trend in fetal deaths per 1,000 births for babies born to Indigenous and non-Indigenous mothers over the 1991–1993 to 2002–2004 periods. Perinatal death rates have not been presented because the data are incomplete for neonatal deaths for some years. Because of the small number of fetal deaths among babies born to Indigenous mothers and the instability in these fetal deaths over time, 3-year moving averages were calculated. As year groupings overlapped, linear regression could not be used when analysing the fetal death trend.

Fetal death rates decreased over time among babies born to Indigenous mothers, from 16 deaths per 1,000 births in 1991–1993 to 11 deaths per 1,000 births in 2002–2004 (Table 6.6). Despite the decrease in fetal deaths among babies born to Indigenous mothers, the rate remained almost double that of the fetal death rate for babies born to non-Indigenous mothers in 2002–2004.

The fetal death rate decreased between 1991–1993 and 2002–2004 for babies born to Indigenous mothers; the rate for babies born to non-Indigenous mothers remained stable over time, at around 6 fetal deaths per 1,000 births.



**Table 6.6: Fetal deaths per 1,000 births to Indigenous and non-Indigenous mothers, 1991–1993 to 2002–2004**

	1991– 1993	1992– 1994	1993– 1995	1994– 1996	1995– 1997	1996– 1998	1997– 1999	1998– 2000	1999– 2001	2000– 2002	2001– 2003	2002– 2004	Average change <sup>(a)</sup>
Indigenous	15.9	15.2	14.8	14.0	14.3	13.3	13.1	12.7	13.1	12.8	12.1	11.4	-0.35*
Non-Indigenous	6.7	6.3	6.4	6.7	6.8	6.7	6.7	6.6	6.7	6.6	6.6	6.9	0.02
<b>Rate ratio</b>	<b>2.4</b>	<b>2.4</b>	<b>2.3</b>	<b>2.1</b>	<b>2.1</b>	<b>2.0</b>	<b>2.0</b>	<b>1.9</b>	<b>2.0</b>	<b>1.9</b>	<b>1.8</b>	<b>1.7</b>	<b>-0.06*</b>

*Note:* Excludes data from Tasmania and the Australian Capital Territory.

*Source:* AIHW NPSU National Perinatal Data Collection 2007.

# Appendix: Technical and explanatory notes

## Technical notes

### Age-standardised rate ratios

In this report, age-standardised rate ratios, using the direct method of standardisation, were reported in Table 2.4 and Table 6.3. The standard population used was all women who gave birth in Australia in 2001.

The direct age-standardised method involves the following steps:

1. Calculate the age-specific rates for each age group. This is done by dividing the number of cases in each specified age group by the corresponding population in that age group. Multiply by 100 to express as a rate per 100.
2. Multiply each age-specific rate by the corresponding standard population in that age group.
3. Sum the results for each age group from Step 2 and divide this sum by the total standard population to obtain the age-standardised rate per 100.

Age-standardised rate ratios were calculated by dividing the Indigenous age-standardised rate by the non-Indigenous age-standardised rate. An age-standardised rate ratio of 1.0 indicates that Indigenous mothers had a rate similar to that of non-Indigenous mothers. An age-standardised rate ratio of 1.2 indicates that Indigenous mothers had a rate that was 20% higher than the rate for non-Indigenous mothers. An age-standardised rate ratio of 0.8 indicates that Indigenous mothers had a rate that was 20% lower than the rate for non-Indigenous mothers.

### Geographical location of usual residence

mapping, the remoteness area data for individual mothers may not be accurate; however, the overall distribution of mothers by geographical areas is considered useful.

Records for mothers not usually resident in Australia and those whose area of usual residence was 'Not stated' were excluded from the tables.

## Socioeconomic status

Data on socioeconomic status are presented throughout Chapters 3 and 4. Socioeconomic Indexes for Areas (SEIFA) provide a summary measure for the socioeconomic conditions within an area. SEIFA were derived from the mother's postcode of usual residence because this was available for all states and territories. The index of advantage/disadvantage, which provides a continuum of advantage to disadvantage, was used. Low values indicate areas of disadvantage and high values indicate areas of advantage.

For 2001–2004, there were 598 records (1.7%) for which an index of advantage/disadvantage was not able to be assigned (Table 3.3). This was because the record was either missing a postcode or did not have a valid postcode.

## Linear regression

Linear regression was used for the analysis of trends in Chapter 6. Linear regression uses the least squares method to calculate a straight line that best fits the data and returns an array that best describes the line.

The simple linear regression line,  $Y = a + bX$ , is normally determined as an estimate from a collection of sample data values consisting of  $X$  values in the scope of the experiment and the corresponding  $Y$  values observed. One common way of estimating the line is the method of least squares. The goal of this method is to create a line that minimizes the summation of the residual error squared. The residual error values are the distances of each sample data point from the resulting best fit line.

Let us use  $e_i$  to represent each residual error squared,  $y_i$  to represent each observed value of  $y$ , and  $\hat{y}_i$  to represent the value of  $Y$  on the estimated line for each  $y_i$ . The method of least squares involves minimising  $e_i = (y_i - \hat{y}_i)^2$ . This is done using partial derivatives, which yield the following formulas for  $a$  ( $y$  intercept estimate) and  $b$  (slope estimate):

$$b = (n \sum x_i y_i - (\sum x_i)(\sum y_i)) / (n \sum x_i^2 - (\sum x_i)^2)$$

$$a = (\sum y_i - b \sum x_i) / n$$

The slope estimate is the average annual change in the data over the period and was used in this report to determine whether there was an increase or decrease in the observed trends in perinatal data over the period 1991–2004. The standard error of the slope estimate was used to determine whether the apparent increases or decreases in the data were statistically significant at the  $p < 0.01$  level.

The formula used to calculate the slope estimate in Microsoft Excel is:

*LINEST* (*known\_y's*, *known\_x's*, *true*) entered as an array formula.





# Glossary

**Aboriginal mothers:** women who gave birth who identified themselves as being of Aboriginal origin. May also include mothers identified as being of both Aboriginal and Torres Strait Islander origin.

**Aboriginal or Torres Strait Islander mothers:** women who gave birth who identified themselves as being of Aboriginal or To

**Mother's length of stay:** number of days between admission date (during the admission resulting in a birth) and separation date (from the hospital where birth occurred). The interval is calculated by subtracting the date of admission from the date of separation.

**Multipara:** pregnant woman who has had at least one previous pregnancy resulting in a live birth or stillbirth.

**Neonatal death:** death of a liveborn baby within 28 days of birth.

**Parity:** number of previous pregnancies resulting in live births or stillbirths, excluding the current pregnancy.

**Perinatal death:** a fetal or neonatal death of at least 20 weeks gestation or at least 400 grams birthweight.

**Plurality:** the number of births resulting from a pregnancy.

**Presentation at birth:** the part of the fetus that appears first at the mouth of the uterus during labour.

**Preterm birth:** birth before 37 completed weeks of gestation.

**Primipara:** pregnant woman who has had no previous pregnancy resulting in a live birth or stillbirth.

**Spontaneous vaginal:** birth without intervention in which the baby's head is the presenting part.

**Stillbirth:** see Fetal death.

**Teenage mother:** mother aged less than 20 years at the birth of her baby.

**Torres Strait Islander mothers:** women who gave birth who identified themselves as being of Torres Strait Islander origin. May also include mothers identified as being of both Torres Strait Islander and Aboriginal origin.

**Vacuum extraction:** assisted birth using a suction cap applied to the baby's head.

**Vaginal breech:** vaginal birth in which the baby's buttocks or lower limbs are the presenting parts.

**Vertex presentation:** The most common and safest birth position where the baby's head is the presenting part.

# References

ABS (Australian Bureau of Statistics) 2004. Experimental estimates and projections, Aboriginal and Torres Strait Islander Australians. ABS cat. no. 3238.0. Canberra: ABS.

ABS & AIHW (Australian Institute of Health and Welfare) 2005. The health and welfare of Australia's Aboriginal and Torres Strait Islander peoples. ABS cat. no. 4704.0. Cat. no. IHW 14. Canberra: ABS & AIHW.

AIHW (Australian Institute of Health and Welfare) 2005. Improving the quality of Indigenous identification in hospital separations data. Cat. no. HSE 101. (Health Services Series no. 25). Canberra: AIHW.

Chan A, Keane R & Robinson J 2001. The contribution of maternal smoking to preterm birth, small for gestational age and low birthweight among Aboriginal and non-Aboriginal births in South Australia. *Medical Journal of Australia* 174(8): 389–93.

Day P, Sullivan EA & Lancaster P 1999. Indigenous mothers and their babies, Australia 1



# List of tables

Table 2.1: Women who gave birth and births by Indigenous status and state and territory, 2001–2004.....	7
Table 2.2: Births to Indigenous mothers by birth status and year, 2001–2004.....	8
Table 2.3: Live births by Indigenous status and state and territory, 2001–2003.....	9
Table 2.4: Summary measures of perinatal health for Indigenous and non-Indigenous mothers and babies, 2001–2004.....	10
Table 3.1: Indigenous mothers by maternal age and state and territory, 2001–2004.....	14
Table 3.2: Indigenous mothers by maternal age and remoteness area of usual residence, 2001–2004.....	15
Table 3.3: Indigenous mothers by maternal age and socioeconomic status, 2001–2004.....	16
Table 3.4: Indigenous mothers by parity and state and territory, 2001–2004.....	17
Table 3.5: Indigenous mothers by maternal age and parity, 2001–2004.....	18
Table 3.6: Indigenous mothers by parity and remoteness area of usual residence, 2001–2004.....	20
Table 3.7: Indigenous mothers by parity and socioeconomic status, 2001–2004.....	21
Table 3.8: Indigenous mothers by tobacco smoking status during pregnancy and state and territory, 2001–2004.....	22
Table 3.9: Indigenous mothers by tobacco smoking status during pregnancy and remoteness area of usual residence, 2001–2004.....	23
Table 3.10: Indigenous mothers by tobacco smoking status during pregnancy and socioeconomic status, 2001–2004.....	24
Table 3.11: Indigenous mothers by tobacco smoking status during pregnancy and maternal age, 2001–2004.....	25
Table 3.12: Indigenous mothers who smoked, by average number of cigarettes per day during the second half of pregnancy and state and territory, 2001–2004.....	26
Table 3.13: Indigenous mothers who smoked during the second half of pregnancy by average number of cigarettes per day and maternal age, 2001–2004.....	27
Table 3.14: Indigenous mothers by onset of labour, type of augmentation or induction and state and territory, 2001–2004.....	28
Table 3.15: Indigenous mothers by presentation of birth and state and territory, 2001–2004.....	29
Table 3.16: Indigenous mothers by method of birth and state and territory, 2001–2004.....	31
Table 3.17: Indigenous mothers by method of birth and remoteness area of usual residence, 2001–2004.....	32
Table 3.18: Indigenous mothers by method of birth and socioeconomic status, 2001–2004.....	33

Table 3.19: Indigenous mothers by actual place of birth and state and territory, 2001–2004 .....	34
Table 3.20: Indigenous mothers who gave birth in hospital by hospital sector, 2001–2004 .....	35
Table 3.21: Indigenous mothers who gave birth in hospital by method of birth and hospital sector, 2001–2004.....	35
Table 3.22: Indigenous mothers who gave birth in hospital <sup>(a)</sup> by postnatal length of stay and maternal age, 2001–2004.....	36
Table 4.1: Live births to Indigenous mothers by sex and state and territory, 2001–2004.....	38
Table 4.2: Preterm births to Indigenous mothers by gestational age and state and territory, 2001–2004.....	39
Table 4.3: Births to Indigenous mothers by gestational age and plurality, 2001–2004.....	41
Table 4.4: Births to Indigenous mothers by gestational age and maternal age, 2001–2004 .....	42
Table 4.5: Live births to Indigenous mothers by birthweight and state and territory, 2001–2004 .....	43
Table 4.6: Live births to Indigenous mothers by birthweight and remoteness area of usual residence, 2001–2004 .....	44
Table 4.7: Live births to Indigenous mothers by birthweight and socioeconomic status, 2001–2004 .....	45
Table 4.8: Live births to Indigenous mothers by birthweight and maternal age, 2001–2004 .....	46
Table 4.9: Mean birthweight (g) of live births to Indigenous mothers by tobacco smoking status during pregnancy and state and territory, 2001–2004.....	47
Table 4.10: Live births to Indigenous mothers by Apgar score at 5 minutes and state and territory, 2001–2004.....	48
Table 4.11: Live births to Indigenous mothers by Apgar score at 5 minutes and remoteness area of usual residence, 2001–2004 .....	48
Table 4.12: Live births to Indigenous mothers in hospital by length of stay and state and territory, 2001–2004.....	50
Table 4.13: Live births to Indigenous mothers in hospital by length of stay and remoteness area of usual residence, 2001–2004 .....	51
Table 4.14: Live births to Indigenous mothers in hospitals, by length of stay and socioeconomic status, 2001–2004 .....	52
Table 4.15: Fetal, neonatal and perinatal deaths of babies born to Indigenous mothers by state and territory, 2001–2004 .....	54
Table 5.1: How the Indigenous status of the mother is obtained, by state and territory, 2007 .....	59
Table 5.2: Whether Indigenous status is commonly checked and/or validated by midwives, by state and territory, 2007.....	60

Table 5.3: Women who gave birth by Indigenous status and state and territory, 1991 to 2004.....	62
Table 5.4: Resident and non-resident women who gave birth in the Australian Capital Territory by Indigenous status, 1993 to 2004 .....	64
Table 6.1: Rates (proportions) and rate ratios of Indigenous and non-Indigenous mothers by maternal age, 1991 to 2004 .....	73
Table 6.2: Rates (proportions) and rate ratios of Indigenous and non-Indigenous first-time mothers by maternal age, 1991 to 2004.....	75
Table 6.3: Age-standardised rate (proportion) and rate ratios of Indigenous and non-Indigenous women who gave birth by caesarean section, 1991 to 2004 .....	77
Table 6.4: Rates (proportion) and rate ratios of low birthweight babies born to Indigenous and non-Indigenous mothers, 1991 to 2004.....	79
Table 6.5: Rates (proportion) and rate ratios of preterm babies born to Indigenous and non-Indigenous mothers, 1991 to 2004.....	81
Table 6.6: Fetal deaths per 1,000 births to Indigenous and non-Indigenous mothers, 1991–1993 to 2002–2004.....	83

# List of figures

Figure 2.1: Age profile of women in the reproductive population by Indigenous status,