

2012 University of New South Wales School Mathematics Competition

Junior Division – Problems and Solutions

Problem 1

Let c be the nested radical

$$c = \sqrt{1 + 2\sqrt{1 + 2\sqrt{1 + 2\sqrt{1 + 2\sqrt{\dots}}}}}$$

converges. Find c .

Solution 1

Note that if

$$c = \sqrt{1 + 2c}$$

More generally it is useful to consider the pattern of numbers in odd

row					1							
row					1	1	1					
row				1	0	1	0	1				
row			1	1	0	1	0	1	1			
row		1	0	0	0	1	0	0	0	1		
row	1	1	1	0	1	1	1	0	1	1	1	
row	1	0	1	0	0	0	1	0	0	0	1	0

Looking at the first four entries in each row starting at row we see that the pattern repeats by construction after row with a 1 in each row

suppose that $b_{k,3} = \frac{1}{2} (k + 1) t$ then

$$\begin{aligned}
 b_{k+1,3} &= b_{k,1} + b_{k,2} + b_{k,3} \\
 &= 1 + k + \frac{1}{2} (k + 1) t
 \end{aligned}$$

straight line directly on up a path of constant gradient they were under a table board after twenty metres and continued walking up the path for another five metres at which point they turned around and noticed that the top of the table board was horizontally with the top of the building. They continued walking up the path a further ten metres where they turned around again and noticed that the top of the building was now visible above the table board. The height of the building is such that it is greater than the height of the

scenario

they return from E to D if the probability of the dice adds up to a number divisible by 3. The probability of returning from E to A is p .

Solution 6

Let P_X denote the probability that the same person stays overnight in town X and let p_{YZ} denote the transition probability that the same person goes from Y to Z .

Then $p_{AB} = 1, p_{BC} = 1, p_{CD} = 1, p_{DE} = 1, p_{ED} = p, p_{EA} = 1 - p$ where $0 \leq p \leq 1$.

We also have $P_A = P_E \times p_{EA}$, $P_B = P_A \times p_{AB}$, $P_C = P_B \times p_{BC}$, $P_D = P_C \times p_{CD} + P_E \times p_{ED}$, $P_E = P_D \times p_{DE}$. Thus $P_A = P_E \times (1 - p)$, $P_B = P_A$, $P_C = P_A$, $P_D = P_A + P_E \times p$, $P_E = P_D$. But $P_A + P_B + P_C + P_D + P_E = 1$ so that $4P_A + (p + 1)P_E = 1$. Finally, substituting P_A from $P_A = P_E \times (1 - p)$ and $4P_A + (p + 1)P_E = 1$ we have $4P_E \times (1 - p) + (p + 1)P_E = 1$ and then $P_E = \frac{1}{-3p + 4}$.

In scenario 1 the sum of the dice is divisible by 3 if the sum is one of 2, 4, 6, 8, 10, 12 so that $p = \frac{1+3+4+3+1}{3} = \frac{12}{3} = 4$ and $P_E = \frac{1}{-3 \times 4 + 4} = \frac{1}{-8} = -\frac{1}{8}$.

In scenario 2 the sum of the dice is divisible by 3 if the sum is one of 3, 6, 9, 12 so that the probability is $p = \frac{2+3+4+1}{3} = \frac{10}{3}$ and $P_E = \frac{1}{-3 \times \frac{10}{3} + 4} = \frac{1}{-26} = -\frac{1}{26}$.

Senior Division – Problems and Solutions

Problem 1

A traveller starts person tours towns A, B, C, D, E and stays overnight in one of the towns. If they stay overnight in town A then the next night they stay in town B . If they stay overnight in town B then the next night they stay in town C . If they stay overnight in town C then the next night they stay in town D . If they stay overnight in town D then the next night they stay in town E . If they stay overnight in town E they return to D for the next night, or go on to town A for the next night. They then continue their tour either from D to E or from A to B etc. At the end of the tour, the probability of finding them in town E on any given night in each of the scenarios below.

Scenario

They return from E to D if the ratio of the distance adds up to a number divisible by two. Otherwise they go on from E to A .

Scenario

They return from E to D if the ratio of the distance adds up to a number divisible by three. Otherwise they go on from E to A .

Solution 1

See solution in the Junior Division part of the book.

Now consider

$$\begin{aligned}
 S &= \sum_{j=0}^{\infty} \frac{j}{n^j} \\
 &= \frac{1}{n} + \frac{2}{n^2} + \frac{3}{n^3} + \frac{4}{n^4} + \frac{5}{n^5} + \dots \\
 &= \frac{1}{n} + \frac{1}{n} \left(\frac{2}{n} + \frac{3}{n^2} + \frac{4}{n^3} + \frac{5}{n^4} + \dots \right) \\
 &= \frac{1}{n} + \frac{1}{n} \left(\frac{1+1}{n} + \frac{1+2}{n^2} + \frac{1+3}{n^3} + \frac{1+4}{n^4} + \dots \right) \\
 &= \frac{1}{n} + \frac{1}{n} \left(\frac{1}{n} + \frac{1}{n^2} + \frac{1}{n^3} + \frac{1}{n^4} + \dots \right) + \frac{1}{n} \left(\frac{1}{n} + \frac{2}{n^2} + \frac{3}{n^3} + \frac{4}{n^4} + \dots \right) \\
 &= \frac{1}{n} + \frac{1}{n} \left(\frac{1}{n} + \frac{1}{n^2} + \frac{1}{n^3} + \frac{1}{n^4} + \dots \right) + \frac{1}{n} S
 \end{aligned}$$

Let us use the known result for the geometric series

$$\frac{1}{n} + \frac{1}{n^2} + \frac{1}{n^3} + \frac{1}{n^4} + \dots = \frac{1}{n-1}$$

we now have


$$S = \frac{1}{n} + \frac{1}{n} \left(\frac{1}{n-1} \right) + \frac{1}{n} S$$

Solution 6

we are given $f(x) = f(x + f(x))$ for a function f so that if we replace x by $x + f(x)$ we have $f(x + f(x)) = f(x + f(x) + f(x + f(x)))$ and if we now use the equality $f(x + f(x)) = f(x)$ we obtain $f(x) = f(x + 2f(x))$. Continuing in this fashion we have $f(x) = f(x + n f(x))$ for a function f and a natural number n .

we consider a proof by contradiction to show that $f(x)$ is a constant function. Suppose that $f(x)$ is not a constant function. Without loss of generality we may assume that there exists $z \in (x, x + f(x))$ such that $f(x) < f(z) < 2f(x)$ and furthermore $f(z) = f(z + n f(z))$ for a natural number n .

Clearly there exists a straight line that separates the point $(z, f(z))$ on the graph from the points $(x, f(x))$ and $(x + f(x), f(x))$. Without loss of generality we suppose that the straight line is given by $y = -\frac{1}{m}x + c$ where m is a positive number. It follows from the continuity of $f(x)$ that there are at least two points $(a, f(a))$ and $(b, f(b))$ with $a \neq b$ that are on the graph and the straight line is its own secant at each of these



us we have $c = a + m f(a)$ and $c = b + m f(b)$ so that $f(c) = f(a + m f(a))$ and $f(c) = f(b + m f(b))$. But $f(a + m f(a)) = f(a)$ and $f(b + m f(b)) = f(b)$ so that $f(c) = f(a) = f(b)$.