MATHEMATICS ENRICHMENT CLUB. Solution Sheet [1](#page-0-0)8, September 15, 2015¹

1. Yes; see gure below.

- 2. We need **k** and **n** such that, $2^{k+1} + \cdots + 2^{k+10} = 1 + 2 + \cdots + n$. Simplifying 2^{k+2} (2^{10} 1) = $n(n + 1)$, which holds for $n = 2^{10}$ 1 and $k = 8$.
- 3. Construct a directed graph with 2000 vertices representing the people, with directed edges connecting the vertices. We want to nd the smallest number of vertices with two edges connecting them. Each vertex has 1000 outgoing edges, so there is a total of 1000 2000 edges. The number of ways we can pair two vertices together is 2000 1999 $2 = 1999$ 1000. So in the best scenario, whereby each vertex of the graph is connected by at least one edge, we still have $2000 - 1000 - 1999 - 1000 = 1000$ edges left over. The extra edges each have to connect vertices that are already connected, hence giving us at least 1000 pairs of friends.
- 4. Three. The Joker can look at the rst three digits of Batman's password and copy them down, then he can then use the fact that the passwords are unique, and deduce

Senior Questions

1. Let

$$
a_n = \frac{r}{n} \frac{\overline{(2n)!}}{n!n^n}.
$$

If we take the natural log on both sides of the above equation, then

$$
\ln(a_n) = \frac{1}{n} \quad \ln \quad \frac{(n+1) \quad (n+2) \quad \dots \quad (n+n)}{n^n}
$$
\n
$$
= \frac{1}{n} \quad \ln \quad \frac{n+1}{n} + \ln \quad \frac{n+2}{n} + \dots + \ln \quad \frac{n+n}{n}
$$
\n
$$
= \frac{1}{n} \ln \quad 1 + \frac{k}{n} \quad \dots
$$

Notice that the RHS of the last displayed equation is a Riemann sum of ln(x) for x 2 [1; 2]. As n gets very large, the Riemann sum for the approximation of $ln(x)$ becomes the exact integral

$$
\begin{array}{c}\nZ_{2} \\
\ln(x) \, dx = 2 \ln 2 \quad 1; \\
1\n\end{array}
$$

 n_1 and n_n

3. Consider the nite sum $S_n = \frac{O_n}{k=2} \cos \frac{\pi}{2^k}$. Multiplying S_n by sin $(\frac{\pi}{2^n})$, then using the trigonometry identity $2 \sin x \cos x = \sin 2x$ yields

sin
$$
\frac{1}{2^n}
$$
 $S_n = \sin \frac{1}{2^n} \cos \frac{1}{2^n} S_{n-1}$
\n $= \frac{1}{2} \sin \frac{1}{2^{n-1}} S_{n-1}$
\n $= \dots$
\n $= \frac{\sin \frac{1}{2}}{2^{n-1}} = \frac{2}{2^n}$:

Therefore, $S_n = \frac{2}{2^n}$ sin $\frac{2}{2^n} = \frac{2}{2^n}$ sin $\frac{2}{2^n}$. But

$$
\lim_{n \to 1} \frac{=2}{\sin(-2)}
$$