## MATHEMATICS ENRICHMENT CLUB. Solution Sheet 12, August 8, 2016

1. Since  $1 + 2 + 3 + \dots + 99 = 4950$  (sum of an arithmetic series), we have

$$n + 2n + 3n + \dots + 99n = n(1 + 2 + 3 + \dots + 99) = 4950n$$


Furthermore, we can factor  $4950 = 5^2 \quad 3^2 \quad 2 \quad 11$ . Therefore, for 4950n to be a perfect square, n must be positive and a multiple of both 11 and 2; the smallest possible n is 22.

2. Since  $1 + 2 + 3 + \cdots + n$  is the sum of an arithmetic series, with common difference 1, we have

$$f(n) = \frac{\frac{n}{2}(2 + (n - 1))}{n} = 1 + \frac{1}{2}(n - 1)$$
:

Therefore  $f(1) + f(2) + \cdots + f(100)$  is sum of an arithmetic series, with initial term 1 and common di erence 0.5. Hence

$$f(1) + f(2) + \dots + f(100) = \frac{100}{2}(2 + 99 \quad 0.5) = 2575$$



3. Let  $P_1; P_2; \ldots; P_n$  be the sides of the polygon, where n is some nonnegative integer. Then for 1 i n, each  $P_i$  is the height of a triangle form by the point P and two adjacent corners of the polygon; for example, as shown in the gure above for the case n = 6. In particular,  $P_i$  hever constitute by  $P_i$  or that  $P_i$  or  $P_i$  is the same for all positions of P.

4. Multiplying both sides of the given equation by  $(x \ 2)(x \ 3)(x \ 5)$  gives

$$x^2$$
  $p = A(x - 3)(x - 5) + B(x - 2)(x - 5) + C(x - 2)(x - 3)$ : (2)

If we substitute x = 2 into (2), then 4 p = 3A. Similarly, substituting x = 3.5 into (2) yields 9 p = 2B and 25 p = 6C. Therefore, we have the system of equations

$$p = 4$$
 3A  
 $p = 9 + 2B$   
 $p = 25$  6C:

The smallest possible p is 7, with A = B = 1 and C = 3.

- 5. We call any 1 cm sides of the lattice an edge, and any point of intersection between edges a vertex. We denote the degree of a vertex by the number of edges incident on that vertex. Consider a vertex with degree 3: since there is an odd number of edges attached to this vertex. Thus, if we can not have multiple threads on an edge, then at least one end of a thread must start at this vertex. In particular, for our 4 cm by 4 cm square lattice, there are 12 vertices with degree 3. Hence, if we only have a total of 40 cm of threads (so that threads can not pass over an edge more than once), then there must be at least 12=2 = 6 pieces of thread to II out the lattice.
  - (a) No possible, too few threads.
  - (b) It is possible.
- 6. Since  $(x + y)^2 = x^2 + 2xy + y$

is 36.

By similar constructions, we can  $\,$  nd the last two digits of  $3^{2016}$ , which is 21. Hence, the required number is 57.

## Senior Questions

- 1. a = 456, b = 546 and c = 1554.
- 2. if  $5a^2 7b^2 = 9$  then 5 does not divide b hence the remainder on dividing b by 5 is 1;2;3 or 4; i.e b = 5c + d, d = 1;2;3 or 4. Therefore

$$b^2 = (25c^2 + 10cd) + d^2$$
;

with  $d^2 = 1/4/9$  or 16. Hence

$$9 = 5a^2$$
  $7b^2 = 5(a^2 35c^2 14cd)$  e;

where e = 7; 28; 63 or 112. In particular,  $5(a^2 35c^2 14cd)$  is equal to 16; 27; 72 or 121, which is impossible. Thus, no such b exists.

3. A convex 8 sided polygon with all angles equal has angles 180  $\frac{360}{8} = 135^{\circ}$ . Hence